arXiv:1509.04918v3 [math.PR] 19 Jul 2016

Convergence Rates for the Degree Distribution in a Dynamic Network Model

Fabian Kück[*] Dominic Schuhmacher[*]

Abstract

In the stochastic network model of Britton and Lindholm [Dynamic random networks in dynamic populations. *Journal of Statistical Physics*, 2010], the number of individuals evolves according to a linear birth and death process with per-capita birth rate λ and per-capita death rate $\mu < \lambda$. A random social index is assigned to each individual at birth, which controls the rate at which connections to other individuals are created. Britton and Lindholm give a somewhat rough proof for the convergence of the degree distribution in this model towards a mixed Poisson distribution. We derive a rate for this convergence giving precise arguments. In order to do so, we deduce the degree distribution at finite time and and derive an approximation result for mixed Poisson distributions to compute an upper bound for the total variation distance to the asymptotic degree distribution. We prove several general results about linear birth and death processes along the way. Most notably, we derive the age distribution of an individual picked uniformly at random at some finite time.

---

[*] Institute for Mathematical Stochastics, University of Göttingen, Goldschmidtstraße 7, 37077 Göttingen, Germany.
Email: fabian.kueck@mathematik.uni-goettingen.de and dominic.schuhmacher@mathematik.uni-goettingen.de

1 Introduction

“Network Science” is a relatively young, rapidly growing research area dealing with complex systems with an underlying graph structure (see e.g. the recent book by van der Hofstad [vdH15]). One of the first random graph models is the well-known Erdős-Rényi model, which is a static model, i.e. it describes a random network at a fixed time. Although this model shows interesting behaviour, further random graph models were needed for the description of real networks since some empirical networks have important properties that are not represented by Erdős-Rényi Graphs. One of these is the power law property of the degree distribution. Preferential attachment models exhibit such a power law behaviour asymptotically and were popularized by Barabási and Albert [BA99]. Peköz, Röllin and Ross [PRR13] established convergence rates for the degree distribution in a discrete-time preferential attachment model in terms of the total variation distance. An alternative to preferential attachment models is the time-continuous random graph model that was introduced by Britton and Lindholm [BL10] and that we consider here. The main difference between this model and the preferential attachment models is that in the former the popularity of each node is determined at birth and stays the same for the whole lifetime whereas the popularity in preferential attachment models is proportional to the degree of the node, which changes over time. Depending on the application either model may be more realistic. In particular it can be shown that the degrees of nodes in the model by Britton and Lindholm can exhibit power laws, such that this model provides an interesting alternative for producing graphs with this property. The fact that time is considered continuously and nodes and connections can not only be created but also destroyed again makes the time-evolution more realistic than in the usual stepwise preferential attachment models.

Let us give a *loop-free* version of the definition of the original *dynamic network model by Britton and Lindholm*; see [BL10] and [BLT11]:
We examine a finite undirected graph without loops that develops over time. The node process $(Y_t)_{t\geq 0}$ is a linear birth and death process with initial value one. Thus each node gives birth at constant rate λ and dies at constant rate μ independently from other nodes. We assume that $(Y_t)_{t\geq 0}$ has right-continuous trajectories. The process $(Y_t)_{t\geq 0}$ and all other random variables that are defined in what follows to describe the dynamic random graph are defined on a common underlying probability space $(\Omega, \mathcal{A}, \mathbb{P})$.

We assume $\lambda > \mu$, so that the node process $(Y_t)_{t\geq 0}$ is a supercritical continuous-time Markov branching process. From standard branching process theory, we obtain that for a random variable W_t with $\mathcal{L}(W_t) = \mathcal{L}(Y_t e^{-t(\lambda-\mu)} | Y_t > 0)$, we have

$$W_t \to W \text{ a.s.},$$

where W is $\mathrm{Exp}(\frac{\lambda}{\lambda-\mu})$ distributed (see e.g. [Har50], page 319).

We equip every node i with a positive random social index S_i, where the S_i are i.i.d. with finite expectation and independent of all other random variables.

This allows us to define the development of the edge set. At birth every node is isolated. During its lifetime and *as long as there is at least one other node*, node i generates and destroys edges according to a birth and death process with constant birth rate αS_i and per-edge death rate β. Here α and β are positive constants. The "second" node of each newly born edge is chosen uniformly at random from the set of all *other* living nodes, and all of the edge processes (including the choices of the second nodes) are independent of each other and all other events.

In addition to the direct destruction of edges in the above process, all edges connected to a certain node are removed when this node dies.

1.1 Remark
The only difference to the definition by Britton and Lindholm is that we do not allow loops because these are not present in most applications. Note that the proofs become slightly simpler if we use the original model by Britton and Lindholm, essentially because times where $Y_t = 1$ need not obtain special treatment. The upper bounds remain largely the same; see also Remark 5.3 for the pure birth case.

Note that we still allow multiple edges. It can be shown that those are negligible in the sense that the probability that a randomly picked node has at least one multiple edge converges to zero at an exponential rate (see Section 6). This allows us to formulate the main result also for the case where we ignore multiple edges (see Corollary 1.4 below).

We refer to the distribution of the number of edges incident to a node picked uniformly at random from all living nodes at time t given the number of nodes is positive as *degree distribution*, and denote it by ν_t. In [BL10], Britton and Lindholm give a somewhat rough proof for the convergence of the degree distribution in the original model towards a mixed Poisson distribution ν. It is the main purpose of this paper to give a rate for the convergence in total variation distance of ν_t to this asymptotic degree distribution ν giving precise arguments. The distribution ν is given by

$$\nu = \mathrm{MixPo}\biggl(\frac{\alpha}{\beta+\mu}\bigl(S + \mathbb{E}(S)\bigr)\bigl(1 - e^{-(\beta+\mu)A}\bigr)\biggr),$$

where $A \sim \mathrm{Exp}(\lambda)$, S has the social index distribution, and A and S are independent. Here MixPo denotes the mixed Poisson distribution. The following result is an immediate consequence of Theorems 5.2 and 5.4 below, which are proved in this paper.

1.2 Theorem
Let $\mathbb{E}(S^2) < \infty$. Then we obtain for the degree distribution ν_t in the Britton–Lindholm Model without loops

(a) if $\mu = 0$, then $d_{TV}(\nu_t, \nu) = O\bigl(\sqrt{t}e^{-\frac{1}{2}\lambda t}\bigr)$ as $t \to \infty$;

(b) if $\mu > 0$, then $d_{TV}(\nu_t, \nu) = O\bigl(t^2 e^{-\frac{1}{6}(\lambda-\mu)t}\bigr)$ as $t \to \infty$.

1.3 Remark
A positive death rate μ leads to a higher variability in the degree distributions for finite t since in

particular the death of a highly connected node (hub) can have a large impact. Thus we would not expect the same rate as in the pure birth case. However, the actual factor in the exponential rate in (b) may be larger than the one stated in the theorem.

Theorem 1.2 has consequences for the case where we ignore multiple edges. Let $\tilde{\nu}_t$ be the distribution of the number of neighbours of a node picked uniformly at random from all living nodes at time t given the number of nodes at time t is positive. We refer to $\tilde{\nu}_t$ shortly as distribution of the number of neighbours. The convergence of this distribution to the asymptotic degree distribution ν is an immediate consequence of the following corollary, which is proved in Section 6.

1.4 Corollary
Let $\mathbb{E}(S^2) < \infty$. For the distribution of the number of neighbours in the Britton–Lindholm Model without loops, we have that $d_{TV}(\tilde{\nu}_t, \nu) = O(t^2 e^{-\frac{1}{6}(\lambda-\mu)t})$ as $t \to \infty$.

1.5 Remark
Britton and Lindholm also introduced a modified model where the "second" node of each newly born edge is not picked uniformly. Instead the probability for each node being the "second" node is proportional to its social index. We expect that one can prove similar results for this modified model in an analogous way. Note that the modification typically leads to a substantially higher ratio of multiple edges, which makes the modified version less interesting for applications.

The rest of the paper is organized as follows. In the second section, we use Poisson approximation to obtain an upper bound for the total variation distance between two mixed Poisson distributions, which we need in Section 5. In Section 3, we consider distributions related to linear birth and death processes since the node population is described by such a process. In particular, we derive the age distribution for a linear birth and death process using results about contour processes from [BPS12]. In Section 4, we derive the degree distribution at finite time by using results about birth and death processes. Finally, in Section 5, we derive rates for the convergence to the asymptotic degree distribution in terms of the total variation distance treating the pure birth case and the general case separately, which leads to Theorem 1.2 above.

2 Upper bound for the total variation distance between two mixed Poisson distributions

Theorem 2.1 in [Yan91] gives us an upper bound for the total variation distance between two Poisson distributions. By conditioning we can generalize this result to mixed Poisson distributions.

2.1 Theorem
Let Λ and M be positive real valued random variables. Then we have for the total variation distance between the mixed Poisson distributions $\mathrm{MixPo}(\Lambda)$ and $\mathrm{MixPo}(\mathrm{M})$:

$$d_{TV}(\mathrm{MixPo}(\Lambda), \mathrm{MixPo}(\mathrm{M})) \leq \mathbb{E}(\min(|\sqrt{\Lambda} - \sqrt{\mathrm{M}}|, |\Lambda - \mathrm{M}|).$$

Proof: Let X and Y be MixPo(Λ) and MixPo(M) distributed, respectively. Then it follows

$$\begin{aligned} d_{TV}(\mathrm{MixPo}(\Lambda), \mathrm{MixPo}(\mathrm{M})) &= \sup_{A\subset\mathbb{N}_0} |\mathbb{P}(X \in A) - \mathbb{P}(Y \in A)| \\ &= \sup_{A\subset\mathbb{N}_0} |\mathbb{E}(\mathbb{P}(X \in A|\Lambda)) - \mathbb{E}(\mathbb{P}(Y \in A|\mathrm{M}))| \\ &\leq \sup_{A\subset\mathbb{N}_0} \mathbb{E}(|\mathbb{P}(X \in A|\Lambda) - \mathbb{P}(Y \in A|\mathrm{M})|) \\ &\leq \mathbb{E}\left(\sup_{A\subset\mathbb{N}_0} |(\mathbb{P}(X \in A|\Lambda)) - (\mathbb{P}(Y \in A|\mathrm{M}))|\right) \\ &= \mathbb{E}(d_{TV}(\mathcal{L}(X|\Lambda), \mathcal{L}(Y|\mathrm{M})) \\ &\leq \mathbb{E}(\min(|\sqrt{\Lambda} - \sqrt{\mathrm{M}}|, |\Lambda - \mathrm{M}|), \end{aligned}$$

where the last line follows from Theorem 2.1 in [Yan91]. □

2.2 Remark
Note that the mixed Poisson distribution depends on the parameter random variable only via its distribution; therefore Theorem 2.1 yields

$$d_{TV}(\mathrm{MixPo}(\Lambda), \mathrm{MixPo}(\mathrm{M})) \leq \inf_{\substack{\tilde{\Lambda}:\tilde{\Lambda}\overset{\mathcal{D}}{=}\Lambda \\ \tilde{\mathrm{M}}:\tilde{\mathrm{M}}\overset{\mathcal{D}}{=}\mathrm{M}}} \mathbb{E}(\min(|\tilde{\Lambda}^{\frac{1}{2}} - \tilde{\mathrm{M}}^{\frac{1}{2}}|, |\tilde{\Lambda} - \tilde{\mathrm{M}}|).$$

3 Distributions related to linear birth and death processes

3.1 Total population size

The node process $(Y_t)_{t\geq 0}$ is a linear birth and death process with birth rate λ, death rate $\mu < \lambda$ and initial value one, i.e. $Y_0 = 1$. Since some results could also be of general interest, we also consider the case where $\mu \geq \lambda > 0$. Firstly, we assume that the birth rate λ and the death rate μ of $(Y_t)_{t\geq 0}$ are not equal. According to (8.15) and (8.46) in [Bai64], the one-dimensional distributions of such a process are given by the following probability mass functions:

$$\begin{aligned} &p_0(t) = \mu\tilde{p}(t), \\ &p_n(t) = (1 - \mu\tilde{p}(t))(1 - \lambda\tilde{p}(t))(\lambda\tilde{p}(t))^{n-1},\ n \geq 1, \end{aligned}$$

where

$$\tilde{p}(t) := \frac{e^{(\lambda-\mu)t} - 1}{\lambda e^{(\lambda-\mu)t} - \mu} = \frac{1}{\lambda}\frac{1 - e^{-(\lambda-\mu)t}}{1 - \frac{\mu}{\lambda}e^{-(\lambda-\mu)t}}.$$

For $\lambda = \mu > 0$, the probability mass function is given by

$$\begin{aligned} p_0(t) &= \frac{\lambda t}{1 + \lambda t}, \\ p_n(t) &= \frac{(\lambda t)^{n-1}}{(1 + \lambda t)^{n+1}},\ n \geq 1, \end{aligned}$$

by (8.53) in [Bai64].

3.1 Remark

Note that $p_0(t)$ is the probability that a linear birth and death process with initial value one goes extinct up to time t. Due to the branching property of a linear birth and death process, we have that $p_0(t)^m$ is the probability that a linear birth and death process with a general initial value m goes extinct up to time t. By taking the limit $t \to \infty$, we obtain that the probability of eventual extinction is $(\mu/\lambda)^m$ for $\lambda \geq \mu$ and 1 otherwise (see e.g. (8.59) in [Bai64]).

By elementary computations using these probability mass functions, we obtain the following proposition (cf. (8.16), (8.17), (8.48), (8.49), (8.54) and (8.55) in [Bai64]).

3.2 Proposition

(i) For $\lambda \neq \mu$, we have

$$\mathbb{E}(Y_t) = e^{(\lambda-\mu)t} \text{ and } \operatorname{Var}(Y_t) = \frac{\lambda+\mu}{\lambda-\mu}\left(e^{2(\lambda-\mu)t} - e^{(\lambda-\mu)t}\right).$$

(ii) For $\lambda = \mu > 0$, we have

$$\mathbb{E}(Y_t) = 1 \text{ and } \operatorname{Var}(Y_t) = 2\lambda t.$$

Now we derive the expected reciprocal of the population size at time t given the population has not gone extinct until time t and state an upper bound:

3.3 Proposition

(i) For $\lambda \neq \mu$, we have

$$\mathbb{E}\left(\frac{1}{Y_t} \,\middle|\, Y_t > 0\right) = \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \lambda} \log\left(\frac{\lambda e^{(\lambda-\mu)t} - \mu}{\lambda-\mu}\right) \leq \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)t\right).$$

(ii) For $\lambda = \mu > 0$ we have,

$$\mathbb{E}\left(\frac{1}{Y_t} \,\middle|\, Y_t > 0\right) = \frac{\log(1+\lambda t)}{\lambda t}.$$

Proof:

(i) We obtain for $\lambda \neq \mu$

$$\begin{aligned}
\mathbb{E}\left(\frac{1}{Y_t} \,\middle|\, Y_t > 0\right) &= \frac{\sum\limits_{n=1}^{\infty} \frac{p_n(t)}{n}}{1 - p_0(t)} \\
&= \sum_{n=1}^{\infty} \frac{1}{n} \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \mu} \left(\frac{\lambda e^{(\lambda-\mu)t} - \lambda}{\lambda e^{(\lambda-\mu)t} - \mu}\right)^{n-1} \\
&= \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \lambda} \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{\lambda e^{(\lambda-\mu)t} - \lambda}{\lambda e^{(\lambda-\mu)t} - \mu}\right)^{n} \\
&= \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \lambda} \left(-\log\left(1 - \frac{\lambda e^{(\lambda-\mu)t} - \lambda}{\lambda e^{(\lambda-\mu)t} - \mu}\right)\right) \\
&= \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \lambda} \log\left(\frac{\lambda e^{(\lambda-\mu)t} - \mu}{\lambda-\mu}\right) \\
&\leq \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \lambda} \left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)t\right).
\end{aligned}$$

(ii) We obtain for $\lambda = \mu > 0$

$$\mathbb{E}\left(\frac{1}{Y_t}\,\middle|\, Y_t > 0\right) = \frac{\sum\limits_{n=1}^{\infty} \frac{p_n(t)}{n}}{1 - p_0(t)} = (1+\lambda t)\sum_{n=1}^{\infty} \frac{1}{n}\frac{(\lambda t)^{n-1}}{(1+\lambda t)^{n+1}} = \frac{1}{\lambda t}\sum_{n=1}^{\infty}\frac{1}{n}\left(\frac{\lambda t}{1+\lambda t}\right)^n = \frac{\log(1+\lambda t)}{\lambda t}.$$

□

In the remaining of this subsection, we only consider the case where $\lambda > \mu$. In this case, we also need an upper bound for $\mathbb{E}(\frac{1}{\sqrt{Y_t}}|Y_t > 0)$, which is given by the following corollary.

3.4 Corollary

For $\lambda > \mu$ and $t \geq \frac{1}{\lambda - \mu}\log(2)$, we have

$$\mathbb{E}\left(\frac{1}{\sqrt{Y_t}}\,\middle|\, Y_t > 0\right) \leq e^{-\frac{1}{2}(\lambda-\mu)t}\sqrt{\frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)t\right)}.$$

Proof: For $\lambda > \mu$ and $t \geq \frac{1}{\lambda-\mu}\log(2)$, it follows from Proposition 3.3:

$$\begin{aligned}
\mathbb{E}\left(\frac{1}{\sqrt{Y_t}}\,\middle|\, Y_t > 0\right) &\leq \sqrt{\mathbb{E}\left(\frac{1}{Y_t}\,\middle|\, Y_t > 0\right)} \\
&\leq \sqrt{\frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t} - \lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)t\right)} \\
&\leq \sqrt{\frac{2(\lambda-\mu)}{\lambda e^{(\lambda-\mu)t}}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)t\right)} \\
&= e^{-\frac{1}{2}(\lambda-\mu)t}\sqrt{\frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)t\right)}.
\end{aligned}$$

□

3.5 Remark

Using Jensen's inequality, we obtain an asymptotic lower bound for the conditional expectation considered in the previous corollary:
By the formula for $p_0(t)$ given at the beginning of this section, it follows that

$$\lim_{t\to\infty} \mathbb{E}(Y_t|Y_t > 0)e^{-(\lambda-\mu)t} = \lim_{t\to\infty}\mathbb{E}(Y_t)e^{-(\lambda-\mu)t}\frac{\lambda}{\lambda-\mu} = \frac{\lambda}{\lambda-\mu} \tag{1}$$

for $\lambda > \mu$, and since the function $x \mapsto \frac{1}{\sqrt{x}}$ is strictly convex on the positive real axis, Jensen's inequality yields

$$\mathbb{E}\left(\frac{1}{\sqrt{Y}_t}\,\middle|\, Y_t > 0\right) > \frac{1}{\sqrt{\mathbb{E}(Y_t|Y_t > 0)}}. \tag{2}$$

Equation (1) implies

$$\lim_{t\to\infty}\frac{1}{\sqrt{\mathbb{E}(Y_t|Y_t>0)}}e^{\frac{1}{2}(\lambda-\mu)t} = \sqrt{\frac{\lambda-\mu}{\lambda}}.$$

Thus we obtain by (2) that $\mathbb{E}(\frac{1}{\sqrt{Y}_t}|Y_t > 0)$ does not converge to zero at a faster rate than $e^{-\frac{1}{2}(\lambda-\mu)t}$ as $t \to \infty$ for $\lambda > \mu$.

3.2 The number of births and deaths

We consider a linear birth and death process $(Y_t)_{t\geq 0}$ with birth rate λ, death rate μ and initial value one as before. For convenience, we assume $\lambda \neq \mu$ in this subsection.

Let B_t be the number of births up to time t. Then $(B_t)_{t\geq 0}$ is in general no Markov process since the number of births depends on the population size. However, the two-dimensional process $(B_t, Y_t)_{t\geq 0}$ is always a Markov process (see Example 6.4 in [CM77]). Let $G(w, z; t) := \mathbb{E}(w^{B_t} z^{Y_t})$ be the joint probability generating function. Then a partial differential equation can be derived for G (see (39) on page 265 in [CM77] or (43) in [Ken49]), which can be used to compute G (see (40) on page 266 in [CM77]). Moreover, the partial differential equation for G leads to the following partial differential equation for the joint cumulant generating function $K(\Theta_w, \Theta_z; t) = \log(G(e^{\Theta_w}, e^{\Theta_z}; t))$ (cf. page 271 in [CM77]):

$$\frac{\partial K}{\partial t} = (\lambda e^{\Theta_w + \Theta_z} - (\lambda + \mu) + \mu e^{-\Theta_z}) \frac{\partial K}{\partial \Theta_z}. \tag{3}$$

Since G is an analytic function, K is also analytic (as a composition of analytic functions). Thus for (Θ_w, Θ_z) from a ball around 0, we can write K in the form

$$K(\Theta_w, \Theta_z; t) = \sum_{i,j=0}^{\infty} \kappa_{ij}(t) \frac{\Theta_w^i \Theta_z^j}{i!j!}, \tag{4}$$

where κ_{ij} denote the corresponding cumulants. By inserting (4) into (3), we obtain

$$\sum_{i,j=0}^{\infty} \frac{d}{dt}\kappa_{ij}(t) \frac{\Theta_w^i \Theta_z^j}{i!j!} = \left(\lambda \sum_{k=0}^{\infty} \frac{\Theta_w^k}{k!} \sum_{l=0}^{\infty} \frac{\Theta_z^l}{l!} - (\lambda + \mu) + \mu \sum_{k=0}^{\infty} \frac{(-\Theta_z)^k}{k!}\right) \sum_{i,j=0}^{\infty} \kappa_{i,j+1}(t) \frac{\Theta_w^i \Theta_z^j}{i!j!}.$$

By equating coefficients, this yields the following differential equations for the cumulants:

$$\frac{d}{dt}\kappa_{01}(t) = (\lambda - \mu)\kappa_{01}(t) \tag{5}$$

$$\frac{1}{2}\frac{d}{dt}\kappa_{02}(t) = (\lambda - \mu)\kappa_{02}(t) + \frac{\lambda + \mu}{2}\kappa_{01}(t) \tag{6}$$

$$\frac{d}{dt}\kappa_{10}(t) = \lambda\kappa_{01}(t) \tag{7}$$

$$\frac{d}{dt}\kappa_{11}(t) = (\lambda - \mu)\kappa_{11}(t) + \lambda\kappa_{02}(t) + \lambda\kappa_{01}(t) \tag{8}$$

$$\frac{1}{2}\frac{d}{dt}\kappa_{20}(t) = \lambda\kappa_{11}(t) + \frac{1}{2}\lambda\kappa_{01}(t). \tag{9}$$

Note that $\kappa_{10}(t) = \mathbb{E}(B_t), \kappa_{01}(t) = \mathbb{E}(Y_t), \kappa_{20}(t) = \operatorname{Var}(B_t), \kappa_{02}(t) = \operatorname{Var}(Y_t)$ and $\kappa_{11}(t) = \operatorname{Cov}(B_t, Y_t)$ (cf. e.g. Subsection 5.1.2 of [CJ10]). Thus from (5) and (6) follows Proposition 3.2. The differential equation (7) yields

$$\mathbb{E}(B_t) = \frac{\lambda}{\lambda - \mu} e^{(\lambda - \mu)t} - \frac{\mu}{\lambda - \mu}$$

if we assume $B_0 = 1$. The differential equation (8) implies

$$\operatorname{Cov}(B_t, Y_t) = \frac{\lambda(\lambda + \mu)}{(\lambda - \mu)^2} e^{2(\lambda - \mu)t} - \frac{2\lambda\mu}{\lambda - \mu} t e^{(\lambda - \mu)t} - \frac{\lambda^2}{(\lambda - \mu)^2} e^{(\lambda - \mu)t}.$$

From (9) follows

$$\frac{d}{dt}\kappa_{20}(t) = \frac{2\lambda^2(\lambda+\mu)}{(\lambda-\mu)^2}e^{2(\lambda-\mu)t} - \frac{4\lambda^2\mu}{\lambda-\mu}te^{(\lambda-\mu)t} - \frac{2\lambda^3}{(\lambda-\mu)^2}e^{(\lambda-\mu)t} + \lambda e^{(\lambda-\mu)t},$$

which implies

$$\operatorname{Var}(B_t) = \frac{\lambda^2(\lambda+\mu)}{(\lambda-\mu)^3}e^{2(\lambda-\mu)t} - \frac{4\lambda^2\mu}{(\lambda-\mu)^2}te^{(\lambda-\mu)t} + \left(\frac{2\lambda^2\mu}{(\lambda-\mu)^3} - \frac{\lambda(\lambda+\mu)}{(\lambda-\mu)^2}\right)e^{(\lambda-\mu)t}.$$

We summarize these results in the following proposition.

3.6 Proposition

If we assume $B_0 = 1$, we have

$$\begin{aligned}
\mathbb{E}(B_t) &= \frac{\lambda}{\lambda-\mu}e^{(\lambda-\mu)t} - \frac{\mu}{\lambda-\mu} \qquad (10)\\
\operatorname{Cov}(B_t, Y_t) &= \frac{\lambda(\lambda+\mu)}{(\lambda-\mu)^2}e^{2(\lambda-\mu)t} - \frac{2\lambda\mu}{\lambda-\mu}te^{(\lambda-\mu)t} - \frac{\lambda^2}{(\lambda-\mu)^2}e^{(\lambda-\mu)t}\\
\operatorname{Var}(B_t) &= \frac{\lambda^2(\lambda+\mu)}{(\lambda-\mu)^3}e^{2(\lambda-\mu)t} - \frac{4\lambda^2\mu}{(\lambda-\mu)^2}te^{(\lambda-\mu)t} + \left(\frac{2\lambda^2\mu}{(\lambda-\mu)^3} - \frac{\lambda(\lambda+\mu)}{(\lambda-\mu)^2}\right)e^{(\lambda-\mu)t}.
\end{aligned}$$

3.7 Remark

The formula (10) can also be computed by using the differential equation

$$\frac{d}{dt}\mathbb{E}(B_t) = \lambda\mathbb{E}(Y_t),$$

which can be derived by conditioning on Y_t (see e.g. Section 2 of [CS12] or proof of Lemma 1 in Section 4 of [BL09]). Let D_t be the number of deaths up to time t. Then we have the analogue differential equation

$$\frac{d}{dt}\mathbb{E}(D_t) = \mu\mathbb{E}(Y_t),$$

which implies

$$\mathbb{E}(D_t) = \frac{\mu}{\lambda-\mu}e^{(\lambda-\mu)t} - \frac{\mu}{\lambda-\mu}$$

if we assume $D_0 = 0$.

3.8 Remark

Note that $\mathbb{E}(Y_t) = \mathbb{E}(B_t) - \mathbb{E}(D_t)$ and $\frac{\mathbb{E}(B_t)-1}{\mathbb{E}(D_t)} = \frac{\lambda}{\mu} = \frac{\lambda}{\lambda+\mu}(\frac{\mu}{\lambda+\mu})^{-1}$ is the ratio of the probabilities of a birth and a death at each event time. Furthermore, $\mathbb{E}(B_t) + \mathbb{E}(D_t) = \frac{\lambda+\mu}{\lambda-\mu}e^{(\lambda-\mu)T} - \frac{2\mu}{\lambda-\mu}$ is the expected number of events up to time t.

3.3 The age distribution

Finally, we consider the age of an individual picked uniformly at random at a fixed time $T > 0$ in the linear birth and death process $(Y_t)_{t\geq 0}$ (given $Y_T > 0$). We briefly call the distribution of this age *the age distribution of* $(Y_t)_{t\geq 0}$ *at time* T (not to be confused with the "age distribution" treated for example in [Har02]). It is known that the age distribution converges to the $\operatorname{Exp}(\lambda)$ distribution as $T \to \infty$; see e.g. [HJV05]. In the pure birth case, the age distribution can be easily derived from known results, such as Theorem 1 in [NR71]. In the general case, the situation is far more complicated. However, we finally obtain the age distribution for any $\mu \geq 0$.

3.3.1 The age distribution in the pure birth case

3.9 Proposition

Let $\mu = 0$. The ages of the individuals at time T that have been born later than the first individual are i.i.d. and their distribution is the truncated exponential distribution with parameter λ. Consequently, the age A of an individual picked uniformly at random from all individuals at time T that were not alive at time 0 satisfies $\mathcal{L}(A) = \mathcal{L}(Z|Z \leq T)$, where Z is $\mathrm{Exp}(\lambda)$ distributed.

Proof: Let $Y_T = n$ be given. From Theorem 1 in [NR71] follows that the $n-1$ positive birth times in the time interval $(0, T]$ are distributed as the order statistics of i.i.d. random variables $\hat{T}_2, \dots, \hat{T}_n$ that have the cumulative distribution function

$$\mathbb{P}(\hat{T}_j \leq u) = \frac{1 - \exp(\lambda u)}{1 - \exp(\lambda T)}.$$

This implies ages $T - \hat{T}_2, \dots, T - \hat{T}_{n-1}$ are i.i.d. and have the cumulative distribution function

$$\mathbb{P}(T - \hat{T}_j \leq u) = 1 - \mathbb{P}(\hat{T}_j \leq T - u) = \frac{1 - \exp(\lambda T) - 1 + \exp(\lambda(T-u))}{1 - \exp(\lambda T)} = \frac{1 - \exp(-\lambda u)}{1 - \exp(-\lambda T)},$$

which is the cumulative distribution function of the truncated exponential distribution with parameter λ. Since this distribution does not depend on n, this conditional distribution is equal to the unconditional one. □

Before we turn to the general case, we give the following corollary.

3.10 Corollary

Let $\mu = 0$ and let $\tilde{A}$ be the age of an (arbitrary) individual at time T and $0 \leq b < T$. Then the conditional distribution of $\tilde{A} - b$ given $\tilde{A} \geq b$ is the exponential distribution with parameter λ truncated at $T - b$.

Proof: Let Z be $\mathrm{Exp}(\lambda)$ distributed. Then for the age $\tilde{A}$ and $z + b \leq T$, Proposition 3.9 implies

$$\begin{aligned}\mathbb{P}(\tilde{A} - b \leq z | \tilde{A} \geq b) = \mathbb{P}(Z \leq z + b | b \leq Z \leq T) = \frac{\mathbb{P}(b \leq Z \leq z + b)}{\mathbb{P}(b \leq Z \leq T)} = \frac{e^{-\lambda b} - e^{-\lambda(z+b)}}{e^{-\lambda b} - e^{-\lambda T}} \\ = \frac{1 - e^{-\lambda z}}{1 - e^{-\lambda(T-b)}} = \mathbb{P}(Z \leq z | Z \leq T - b)\end{aligned}$$

and $z \mapsto \mathbb{P}(Z \leq z | Z \leq T - b)$ is the cumulative distribution function of the exponential distribution with parameter λ truncated at $T - b$. □

3.11 Remark

From the proof of Proposition 3.9, we know that we obtain the same distribution in Proposition 3.9 and Corollary 3.10 if we condition on the population size Y_T. Note that the age distribution of an individual picked uniformly at random conditioned on Y_T follows immediately. The unconditional age distribution can be easily deduced from the conditional one by applying Proposition 3.3. The result is omitted here because the cumulative distribution function of the age distribution for arbitrary $\mu \geq 0$ is stated in Corollary 3.13 below.

3.3.2 The age distribution in the general case

We consider the general case with birth rate $\lambda > 0$ and death rate μ. For convenience, we assume $\mu > 0$ although all results also hold for $\mu = 0$.

The following theorem gives us the age distribution at some time T conditioned on the population size.

3.12 Theorem

Let F_{y_T} denote the cumulative distribution function of the age of an individual picked uniformly at random at time T given $Y_T = y_T$ for some $y_T > 0$. Then F_{y_T} is given by

$$F_{y_T}(t) = \frac{y_T - 1}{y_T}\left(1 - \frac{e^{-\lambda t} - e^{-(\lambda-\mu)T}e^{-\mu t}}{1 - e^{-(\lambda-\mu)T}}\right) + \frac{1}{y_T}\left(\frac{\lambda(1-e^{-\mu t}) - \mu(1-e^{-\lambda t})}{\lambda - \mu}\mathbb{1}_{\{t<T\}} + \mathbb{1}_{\{t=T\}}\right)$$

for $t \in [0, T]$ if $\lambda \neq \mu$ and by

$$F_{y_T}(t) = \frac{y_T - 1}{y_T}\left(1 - \frac{e^{-\lambda t}(T-t)}{T}\right) + \frac{1}{y_T}\left(1 - e^{-\lambda t}(1+\lambda t))\mathbb{1}_{\{t<T\}} + \mathbb{1}_{\{t=T\}}\right)$$

for $t \in [0, T]$ if $\lambda = \mu > 0$.

Before we prove this theorem, we state two corollaries. The first one reveals the unconditional age distribution of an individual picked uniformly at random.

3.13 Corollary

The cumulative distribution function F of the age of an individual picked uniformly at random at time T is given by

$$\begin{aligned} F(t) =& \left(1 - \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)T} - \lambda}\log\left(\frac{\lambda e^{(\lambda-\mu)T} - \mu}{\lambda-\mu}\right)\right)\left(1 - \frac{e^{-\lambda t} - e^{-(\lambda-\mu)T}e^{-\mu t}}{1 - e^{-(\lambda-\mu)T}}\right) \\ &+ \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)T} - \lambda}\log\left(\frac{\lambda e^{(\lambda-\mu)T} - \mu}{\lambda-\mu}\right)\left(\frac{\lambda(1-e^{-\mu t}) - \mu(1-e^{-\lambda t})}{\lambda-\mu}\mathbb{1}_{\{t<T\}} + \mathbb{1}_{\{t=T\}}\right) \end{aligned}$$

for $t \in [0, T]$ if $\lambda \neq \mu$ and by

$$\begin{aligned} F(t) = &\left(1 - \frac{\log(1+\lambda t)}{\lambda t}\right)\left(1 - \frac{e^{-\lambda t}(T-t)}{T}\right) \\ &+ \frac{\log(1+\lambda t)}{\lambda t}\left((1 - e^{-\lambda t}(1+\lambda t))\mathbb{1}_{\{t<T\}} + \mathbb{1}_{\{t=T\}}\right) \end{aligned}$$

for $t \in [0, T]$ if $\lambda = \mu > 0$.

Proof: The expression for the cumulative distribution function follows directly from Theorem 3.12 and Proposition 3.3. □

Note that the cumulative distribution function in Corollary 3.13 converges to the cumulative distribution function of the Exp(λ) distribution as $T \to \infty$. The following corollary states that the age distribution converges exponentially fast to the Exp(λ) distribution in a certain sense for $\lambda > \mu$.

3.14 Corollary
Let $\lambda > \mu$ and let A denote the age of an individual picked uniformly at random at time T in the linear birth and death process $(Y_t)_{t\geq})$ (given $Y_t > 0$). Then there exists a random variable Z with $\mathcal{L}(Z|Y_T > 0) = \mathrm{Exp}(\lambda)$ such that

$$\mathbb{E}\left(\left|e^{-cA} - e^{-cZ}\right| \,\middle|\, Y_T > 0\right) \leq \frac{\lambda}{c+\lambda} \frac{1}{e^{(\lambda-\mu)T} - 1} + \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)T} - \lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)T\right)$$

for any $c > 0$.

In order to prove Theorem 3.12 and Corollary 3.14, we use results from [BPS12] about contour processes (also known as exploration processes), which we introduce now.

Bijection between piecewise-linear functions and trees

3.15 Definition
We consider piecewise-linear functions $h : [0, b] \to [0, \infty)$ for $0 < b < \infty$ with the following properties:

- The derivative is either 1 or -1 at all points where it exists.[1]
- $h(x) > 0$ if and only if $x \in (0, b)$
- All non-zero local minima are at distinct heights.

We identify each such function $h : [0, b] \to [0, \infty)$ with the function $\tilde{h} : [0, \infty) \to [0, \infty)$ that fulfils

$$\tilde{h}(x) := \begin{cases} h(x) & \text{if } 0 \leq x \leq b \\ 0 & \text{if } x > b. \end{cases}$$

Let $\mathcal{H}$ denote the set of all such functions.

Now we introduce a bijection Φ between $\mathcal{H}$ and the set $\mathcal{T}$ of finite rooted binary trees. If not stated otherwise, we draw trees in a left-aligned way in this subsection, i.e. each inner node has a vertical line going in and out, and one horizontal line going to the right (see Figure 1). Horizontal lines are for constructive purposes only, whereas the vertical line segments represent actual edges and the corresponding edge lengths in the tree. Further below we will identify such trees with realizations of birth and death processes that also include information about the lifetimes of the individuals represented by the lengths of the vertical lines.

Starting from a piecewise linear function $h \in \mathcal{H}$, we define $\Phi(h)$ by drawing a tree under the graph of h as follows (see also Figure 1):
Let $(x_{\max,1}, h_{\max,1})$ be the leftmost local maximum of h. Then we draw a (vertical) line from $(x_{\max,1}, 0)$ to $(x_{\max,1}, h_{\max,1})$. Let $(x_{\max,2}, h_{\max,2})$ be the second leftmost local maximum and $(x_{min,1}, h_{min,1})$ be the leftmost non-zero local minimum if they exist. In this case we add a (vertical) line from $(x_{\max,2}, h_{min,1})$ to $(x_{\max,2}, h_{\max,2})$ and a (horizontal) line from $(x_{\max,2}, h_{min,1})$ connecting the vertical

---

[1] Note that the formulation in [BPS12] is slightly more general. However, we do not need this more general setting for our purposes.



Figure 1: A piecewise linear function and the corresponding (left-aligned) tree

line horizontally to the rest of the tree. We continue this procedure until we have explored all non-zero local extrema (see Figure 1).

It is clear that Φ is a bijection since we obtain the inverse of Φ by assigning to any (left-aligned) tree, where the horizontal position of the root and the horizontal line lengths have been suitably adapted, the function that starts at $(0,0)$ and “turns” at every leaf and every time it hits a horizontal line until it hits the x-axis again.

Probability laws of the contour process and the corresponding random tree

Now we define probability measures on $\mathcal{H}$ and $\mathcal{T}$. Let for the moment $0 < \lambda \leq \mu$. We define the contour process as a stochastic process $(H_x)_{x \geq 0}$ whose trajectories belong to $\mathcal{H}$. Let $(U_k)_{k \in \mathbb{N}}$ and $(V_k)_{k \in \mathbb{N}}$ be two mutually independent sequences of independent, identically exponentially distributed random variables with parameters μ and λ, respectively. Let $Z_k = U_k - V_k$ for all k and let $P_{\lambda,\mu}$ be the law of the random element of $\mathcal{H}$ with the following properties. Starting with a first local minimum at $(0,0)$, the heights of the first local maximum and the second local minimum are U_1 and $(Z_1)^+$, respectively. The process is stopped whenever it returns to zero. If $(Z_1)^+ > 0$, the heights of the second local maximum and the third local minimum are $U_2 + Z_1$ and $(Z_1 + Z_2)^+$, respectively. If $(Z_1 + Z_2)^+ > 0$, we continue in the same way (see Figure 2).



Figure 2: The contour process

It can be shown that this process returns to zero almost surely if $0 < \lambda \leq \mu$ (see Section 2 of [BPS12]).

In Theorem 3.16 below, we relate $P_{\lambda,\mu}$ to the law of a linear birth death process with the same parameters. Since we are also interested in the supercritical case, we also define the law of a random element of $\mathcal{H}$ that can be related to the law of a supercritical linear birth and death process with birth rate λ and death rate $\mu < \lambda$ now. Since the contour process defined above does not return to zero almost surely if $\lambda > \mu$, we need to modify the construction. Therefore, we take the process from above and add a deflection at T: The j-th time the process reaches T, it stops and goes down with slope -1 for a time $\min(T, \hat{Z}_j)$ where $\hat{Z}_j \sim \text{Exp}(\lambda)$ independent (also from all other random variables). It can be shown that this deflected process returns to zero almost surely (see Section 2 of [BPS12]). Let $P_{\lambda,\mu,T}$ denote the law of the deflected process. Note that $P_{\lambda,\mu,\infty} = P_{\lambda,\mu}$.

Let $Q_{\lambda,\mu,T}$ be the law of the random tree corresponding to a linear birth and death process with parameters λ and μ killed at time T, i.e. all individuals alive at time $T-$ are killed at time T. From a realization of such a birth and death process, including information about the lifetimes of the individuals, draw the corresponding random tree as follows. Draw the lifetime of the individual that is alive at time 0 as the leftmost vertical line. For each of its direct offspring attach a horizontal line and draw their lifetimes as further vertical lines. Starting with the latest offspring and keeping the lower horizontal lines long enough will keep the tree planar. Continue in the same way for the offspring of each offspring and so on. Note that the lifetimes of the individuals correspond to the lengths of the vertical lines. Obviously, the ages of the individuals at time T in a linear birth and death process are just obtained by the total lifetimes of the individuals alive at time $T-$ in the corresponding killed tree (see Figure 3).

We allow $T = \infty$, in which case the birth and death process is not killed.

The following theorem reveals the simple relationship between $Q_{\lambda,\mu,T}$ and $P_{\lambda,\mu,T}$.

3.16 Theorem

For any $\lambda, \mu > 0$ and $T \in (0, \infty)$, and also for $T = \infty$ if $\lambda \leq \mu$, we have

$$Q_{\lambda,\mu,T} = P_{\lambda,\mu,T}\Phi^{-1}.$$

Proof: [BPS12], Theorem 3.1. See the proof of that theorem to verify the precise construction of the left-aligned tree based on the birth and death process. ☐

Proof of Theorem 3.12 and Corollary 3.14

We first prove that we obtain the same distribution if ages are read off from the tree under the vertically mirrored contour process $(H_{\tau-t})_{0\leq t\leq\tau}$, where $\tau = \inf\{x > 0;\, H_x = 0\}$ (see Figure 4). To see this, denote by $\Psi\colon \mathcal{T} \to \mathcal{T}$ the corresponding transformation on the set of trees. Let $D \sim Q_{\lambda,\mu,T}$ be the random tree obtained from the above birth and death process and denote by $\Theta(d)$ the finite set of trees that have the same vertex positions in the vertical direction as a tree $d \in \mathcal{T}$, but not necessarily the same phylogeny. Note that $\Theta(\Psi(d)) = \Theta(d)$ for every $d \in \mathcal{T}$. Given $\Theta(D)$, the random tree D is uniformly distributed on $\Theta(D)$, and so is $\Psi(D)$ because, restricted to the finite set $\Theta(D)$, the map Ψ is just a permutation. Hence the distributions of $\Psi(D)$ and D are equal by integration of the conditional distributions given $\Theta(D)$.



Figure 3: The deflected contour process and the corresponding tree representing a linear birth and death process up to time T



Figure 4: The mirrored deflected contour process (the realization corresponds to Figure 3) and its (left-aligned) tree

Since it is more customary to draw stochastic processes from left to right, we mirror the situation in Figure 4, which gives us the original contour process (H_x) again, enveloping a right-aligned tree (constructed in the analogous way as the left-aligned tree, but from right to left; see Figure 5). Of course the ages still correspond to vertical line lengths.

Obviously, the local maxima of the contour process at height T correspond to the living individuals in the linear birth and death process at time T, which we assume to be numbered from left to right in the right-aligned tree here. In total, there are B_T local maxima (at any height) corresponding to the B_T individuals that have been born up to time T. Let these maxima be numbered from left to right and let $I_1 < \ldots < I_{Y_T}$ be the indices of the maxima at height T and $I_k = \infty$ for $k > Y_T$. Then $V_{I_1}, \ldots, V_{I_{Y_T - 1}}, V_{I_{Y_T}} \wedge T$ are the ages we are interested in, where $(V_k)_{k \in \mathbb{N}}$ is the sequence of random variables from the definition of the contour process, i.e. $(V_k)_{k \in \mathbb{N}}$ are i.i.d. $\operatorname{Exp}(\lambda)$ distributed. Let $0 = W_1 < \ldots < W_{2B_T+1} = \tau$ be the positions of all local extrema of $(H_x)_{0 \leq x \leq \tau}$. Note that $W_{2I_1}, \ldots, W_{2I_{Y_T}}$ are the positions of the maxima at height T. For convenience, we define $W_k = W_{2B_T+1} = \tau$ for all $k > 2B_T$ and $W_\infty = \lim\limits_{k \to \infty} W_k = \tau$.

By the above construction of the contour process, the process $(H_{W_k}, (-1)^k)_{k \in \mathbb{N}}$ is a Markov chain



Figure 5: The deflected contour process and its right-aligned tree

on $[0,T] \times \{-1,1\}$ with initial value $(0,-1)$ at time 1 and transition probabilities

$$\begin{aligned}&\mathbb{P}((H_{W_{k+1}},(-1)^{k+1}) \in B_1 \times B_2 | (H_{W_k},(-1)^k) = (z_1,z_2)) \\ &\quad = \begin{cases} \Big(\delta_T(B_1)e^{-\mu(T-z_1)} + \mu \int\limits_{B_1\cap[z_1,T)} e^{-\mu(u-z_1)}du\Big)\delta_1(B_2) & \text{if } z_2 = -1 \\ \Big(\delta_0(B_1)e^{-\lambda z_1} + \lambda \int\limits_{B_1\cap(0,z_1]} e^{-\lambda(z_1-u)}du\Big)\delta_{-1}(B_2) & \text{if } z_2 = 1, \end{cases}\end{aligned}$$

where B_1 is a Borel subset of $[0,T]$, $B_2 \subset \{0,1\}$ and $(z_1,z_2) \in [0,T] \times \{-1,1\}$. Note that the second component of $(H_{W_k},(-1)^k)_{k\in\mathbb{N}}$ tells us if $(H_x)_{0\leq x\leq\tau}$ has a minimum (if it is -1) or a maximum (if it is 1) at W_k. Note further that this Markov chain determines $(H_x)_{x\geq 0}$ completely.

We define the sequence $(\xi_k)_{k\in\mathbb{N}}$ of hitting times in $(T,1)$ recursively by

$$\xi_1 = \inf\{k \in \mathbb{N} : (H_{W_k},(-1)^k) = (T,1)\} \text{ and } \xi_{l+1} = \inf\{k > \xi_l : (H_{W_k},(-1)^k) = (T,1)\}$$

for all $l \in \mathbb{N}$, where $\inf\emptyset = \infty$. Note that each ξ_l is a stopping time and that $\max\{l : \xi_l < \infty\} = Y_T$.

Moreover, we define the "horizontally mirrored" excursions

$$E^{(l)} = (E_k^{(l)})_{0\leq k\leq\xi_{l+1}-\xi_l} = (T - H_{W_k})_{\xi_l\leq k\leq\xi_{l+1}}$$

for $l \in \mathbb{N}$, where we set $\infty - \infty = 0$.

Since $\mathbb{P}((H_{W_{\xi_l}},(-1)^{\xi_l}) = (T,1)|\xi_l < \infty) = 1$, the strong Markov property and the fact that the second component of the process is deterministic imply that $(H_{W_k})_{0\leq k\leq\xi_l}$ and $(H_{W_k})_{\xi_l\leq k}$ are independent given $\xi_l < \infty$ and that $(H_{W_k})_{\xi_l\leq k}$ has the same distribution given $\xi_l < \infty$ for any $l \in \mathbb{N}$. As a consequence, we have for any $l \in \mathbb{N}$ that $(H_{W_k})_{0\leq k\leq\xi_1}, E^{(1)}, E^{(2)}, \dots, E^{(l-1)}$ are independent of $(E^{(l)},(\xi_r)_{r\geq l+1})$ given $\xi_l < \infty$ and hence that $(H_{W_k})_{0\leq k\leq\xi_1}, E^{(1)}, E^{(2)}, \dots, E^{(l-1)}$ are independent of $E^{(l)}$ given $\xi_l < \infty$ and given any sub-σ-algebra of $\sigma(\xi_r; r \geq l+1)$. This implies that $(H_{W_k})_{0\leq k\leq\xi_1}, E^{(1)}, E^{(2)}, \dots, E^{(l-1)}, E^{(l)}$ are independent given $\xi_l < \infty$ and given any sub-σ-algebra of $\sigma(\xi_r; r \geq l+1)$ for any $l \in \mathbb{N}$. Note that for any $l \in \mathbb{N}$, the first $l-1$ "horizontally mirrored" excursions $E^{(1)}, E^{(2)}, \dots, E^{(l-1)}$ all have the same distribution under this conditioning because $E^{(1)}, E^{(2)}, \dots, E^{(l-1)}$ all have the same distribution given $\xi_l < \infty$ and $E^{(1)}, E^{(2)}, \dots, E^{(l-1)}$ are independent of $(\xi_r)_{r\geq l+1}$ given $\xi_l < \infty$.

Since $\{Y_T = y_T\} = \{\xi_{y_T} < \infty, \xi_{y_T+1} = \infty\}$ we obtain that, given $Y_T = y_T$, the processes $(H_{W_k})_{0\le k\le\xi_1}, E^{(1)}, E^{(2)}, \dots, E^{(y_T-1)}, E^{(y_T)}$ are independent and $E^{(1)}, E^{(2)}, \dots, E^{(y_T-1)}$ are identically distributed. By having a closer look at the strong Markov property used above, it is seen that neither the distribution of $E^{(1)}$ given $Y_T = y_T \ge 2$ nor the distribution of $E^{(y_T)}$ given $Y_T = y_T \ge 1$ depend on the concrete value y_T. Note that on $\{Y_T = y_T\}$, we have $E_1^{(1)} = V_{I_1}, \dots, E_1^{(y_T-1)} = V_{I_{y_T-1}}$, and $E_1^{(y_T)} = V_{I_{y_T}} \wedge T$, so these are the ages we are interested in. The conditional distributions of $E_1^{(1)}$ and $E_1^{(y_T)}$ given $Y_T = y_T$ can be easily rediscovered as conditional distributions in another contour process, which is what we look at next.

Let $(\tilde{H}_x)_{x\ge0} \sim P_{\mu,\lambda,T}$. By Theorem 3.16, $(\tilde{H}_x)_{x\ge0}$ is the contour process corresponding to a linear birth and death process with *birth rate* μ and *death rate* λ that is killed at T. Let the positions $(\tilde{W}_k)_k$ of the extrema of $(\tilde{H}_x)_{x\ge0}$ be defined analogously to $(W_k)_k$.

Both $(\tilde{H}_{\tilde{W}_{k+1}})_{k\in\mathbb{N}_0}$ and $(E_k^{(l)})_{k\in\mathbb{N}_0}$ for arbitrary $l \in \{1, \dots, Y_T\}$ start in 0 and alternate between independently adding $\mathrm{Exp}(\lambda)$ and subtracting $\mathrm{Exp}(\mu)$ random variables at least until 0 or T is crossed. Thus until this happens, they have the same distribution. We have $\{Y_T \ge 2\} = \{E^{(1)}_{\xi_2-\xi_1} = 0\} = \{\max_k E_k^{(1)} < T\}$, and therefore for any $y_T \ge 2$ that, since the distribution of $E_1^{(1)}$ does not depend on the concrete value of y_T,

$$\mathcal{L}(E_1^{(1)} \,|\, Y_T = y_T) = \mathcal{L}\Big(\tilde{H}_{\tilde{W}_2} \,\Big|\, \max_k \tilde{H}_{\tilde{W}_k} < T\Big) = \mathcal{L}\Big(\tilde{H}_{\tilde{W}_2} \,\Big|\, \max_{x\ge0} \tilde{H}_x < T\Big). \tag{11}$$

On the other hand we have $\{Y_T \ge 1\} = \{Y_T \ge 1, \max_k E_k^{(Y_T)} = T\}$ (in fact $\xi_{Y_T+1} = \infty$ and $E^{(Y_T)}$ is eventually absorbed in T). Therefore for any $y_T \ge 1$, since the distribution of $E_1^{(y_T)}$ does not depend on the concrete value of y_T,

$$\mathcal{L}(E_1^{(y_T)} \,|\, Y_T = y_T) = \mathcal{L}\Big(\tilde{H}_{\tilde{W}_2} \,\Big|\, \max_k \tilde{H}_{\tilde{W}_k} = T\Big) = \mathcal{L}\Big(\tilde{H}_{\tilde{W}_2} \,\Big|\, \max_{x\ge0} \tilde{H}_x = T\Big). \tag{12}$$

In other words, we have identified the desired age distributions at time T given $Y_T = y_T$ as the distribution of the lifetime of the first individual in a linear birth and death process with *birth rate* μ and *death rate* λ conditioned on extinction of the process by time T (Equation (11), first $y_T - 1$ individuals) or conditioned on survival of the process up to time T (Equation (12), last individual, lifetime measured up to time T).

We first compute the age distribution of the first $y_T - 1$ individuals given $Y_T = y_T \ge 2$. By Bayes' Theorem $\mathcal{L}(\tilde{H}_{\tilde{W}_2} \,|\, \max_{x\ge0} \tilde{H}_x < T)$ has a density

$$f_*(t) \propto \lambda e^{-\lambda t}\mathbb{P}\Big(\max_{x\ge0} \tilde{H}_x < T \,\Big|\, \tilde{H}_{\tilde{W}_2} = t\Big), \tag{13}$$

where $t < T$. By Theorem 3.16, we have to compute the probability that the linear birth and death process with birth rate μ and death rate λ goes extinct before time T given the lifetime of the first individual is t. Note that the birth times of the offspring of the first individual form a Poisson process with rate μ (up to time t). By conditioning on their number (up to time t) and their birth times, and by the formula for the extinction probability from Subsection 3.1, we obtain that this probability is equal to

$$\begin{aligned}
&\sum_{k=0}^{\infty}\frac{(\mu t)^k}{k!}e^{-\mu t}\frac{1}{t^k}\int_0^t\cdots\int_0^t\frac{\lambda}{\mu}\frac{1-e^{(\lambda-\mu)(T-t_1)}}{1-\frac{\lambda}{\mu}e^{(\lambda-\mu)(T-t_1)}}\cdots\frac{\lambda}{\mu}\frac{1-e^{(\lambda-\mu)(T-t_k)}}{1-\frac{\lambda}{\mu}e^{(\lambda-\mu)(T-t_k)}}dt_1\dots dt_k\\
&=\sum_{k=0}^{\infty}\frac{1}{k!}e^{-\mu t}\left(\int_0^t\lambda\frac{e^{-(\lambda-\mu)(T-u)}-1}{e^{-(\lambda-\mu)(T-u)}-\frac{\lambda}{\mu}}du\right)^k\\
&=\sum_{k=0}^{\infty}\frac{1}{k!}e^{-\mu t}\Big(\log(\lambda e^{(\lambda-\mu)(T-t)}-\mu)+\lambda t-\log(\lambda e^{(\lambda-\mu)T}-\mu)\Big)^k\\
&=\frac{\lambda e^{(\lambda-\mu)(T-t)}-\mu}{\lambda e^{(\lambda-\mu)T}-\mu}e^{(\lambda-\mu)t}
\end{aligned}\tag{14}$$

for $t < T$ if $\lambda \neq \mu$. For $\lambda = \mu > 0$, we see analogously that this probability is equal to

$$\begin{aligned}
&\sum_{k=0}^{\infty}\frac{(\lambda t)^k}{k!}e^{-\lambda t}\frac{1}{t^k}\int_0^t\cdots\int_0^t\frac{\lambda(T-t_1}{1+\lambda(T-t_1)}\cdots\frac{\lambda(T-t_k}{1+\lambda(T-t_k)}dt_1\dots dt_k\\
&=\sum_{k=0}^{\infty}\frac{1}{k!}e^{-\lambda t}\left(\int_0^t\lambda\frac{\lambda(T-u)}{1+\lambda(T-u)}du\right)^k\\
&=\sum_{k=0}^{\infty}\frac{1}{k!}e^{-\lambda t}(\log(1+\lambda(T-t))-\log(1+\lambda T)+\lambda T)^k\\
&=\frac{1+\lambda(T-t)}{1+\lambda T}
\end{aligned}\tag{15}$$

Now we can easily compute the normalizing constant of the density f_* in (13): For $\lambda \neq \mu$, we obtain

$$\begin{aligned}
\int_0^T\lambda e^{-\lambda t}(\lambda e^{(\lambda-\mu)(T-t)}-\mu)e^{(\lambda-\mu)t}dt&=\int_0^T(\lambda e^{(\lambda-\mu)T}e^{-\lambda t}-\mu e^{-\mu t})dt\\
&=e^{(\lambda-\mu)T}(1-e^{-\lambda T})-(1-e^{-\mu T})\\
&=e^{(\lambda-\mu)T}-1
\end{aligned}$$

and for $\lambda = \mu > 0$, we have

$$\begin{aligned}
\int_0^T\lambda e^{-\lambda t}(1+\lambda(T-t))dt&=(1-e^{-\lambda T})(1+\lambda T)-\int_0^T\lambda^2e^{-\lambda t}t\,dt\\
&=(1-e^{-\lambda T})(1+\lambda T)-(1-e^{-\lambda T}(\lambda T+1))\\
&=\lambda T.
\end{aligned}$$

Thus the density f_* is given by

$$f_*(t)=\frac{\lambda e^{(\lambda-\mu)T}e^{-\lambda t}-\mu e^{-\mu t}}{e^{(\lambda-\mu)T}-1}=\frac{\lambda e^{-\lambda t}-\mu e^{-(\lambda-\mu)T}e^{-\mu t}}{1-e^{-(\lambda-\mu)T}}$$

for $t \in [0,T]$ if $\lambda \neq \mu$ and by

$$f_*(t) = \frac{\lambda e^{-\lambda t}(1+\lambda(T-t))}{\lambda T} = \frac{e^{-\lambda t}(1+\lambda(T-t))}{T}$$

for $t \in [0,T]$ if $\lambda = \mu > 0$. By integration, we obtain that the cumulative distribution function F_* for the age of any of the first $y_T - 1$ individuals given $Y_T = y_T$ takes the form

$$F_*(t) = 1 - \frac{e^{-\lambda t} - e^{-(\lambda-\mu)T}e^{-\mu t}}{1-e^{-(\lambda-\mu)T}} \tag{16}$$

and

$$F_*(t) = 1 - \frac{e^{-\lambda t}(T-t)}{T}$$

for $t \in [0,T]$, respectively. By multiplying the probability that, given $Y_T = y_T$, the randomly picked individual is one of the first $y_T - 1$ individuals, we obtain the first summands in Theorem 3.12.

3.17 Remark
In particular, we see that

$$1 - F_*(t) = \frac{e^{-\lambda t} - e^{-(\lambda-\mu)T}e^{-\mu t}}{1-e^{-(\lambda-\mu)T}} \leq \frac{e^{-\lambda t}(1-e^{-(\lambda-\mu)T})}{1-e^{-(\lambda-\mu)T}} = e^{-\lambda t}$$

for $t \in [0,T]$ if $\lambda \neq \mu$ and

$$1 - F_*(t) = \frac{e^{-\lambda t}(T-t)}{T} \leq e^{-\lambda t}$$

for $t \in [0,T]$ if $\lambda = \mu$. Thus, given $Y_T = y_T$, the age distribution of the first $y_T - 1$ individuals is stochastically dominated by the $\mathrm{Exp}(\lambda)$ distribution.

Before we determine the distribution of the last individual in a similar way, we prove Corollary 3.14. We can explicitly define the random variable Z in the statement of this corollary. Given $Y_T = y_T$, let J_T be uniformly distributed on $\{1, ..., y_T\}$, independent of all other random variables, and let A be the age of individual J_T, where the individuals alive at time T are still numbered from left to right in the right-aligned tree.

Let $\tilde{Z}$ be an arbitrary $\mathrm{Exp}(\lambda)$ distributed random variable that is independent of everything else and let F_∞ denote the cumulative distribution function of the $\mathrm{Exp}(\lambda)$ distribution. Set

$$Z(\omega) := \mathbb{1}_{\{J_T(\omega)<Y_T(\omega)\}}F_\infty^{-1}(F_*(A(\omega)) + \mathbb{1}_{\{J_T(\omega)=Y_T(\omega)\}}\tilde{Z}.$$

Then Z is $\mathrm{Exp}(\lambda)$ distributed given $Y_T > 0$ because conditionally on $Y_T = y_T > 0$ it is $\mathrm{Exp}(\lambda)$ distributed both given $J_T = y_T$ and given $J_T < y_T$. The latter holds because F_* is a continuous function. We have for $c > 0$

$$\begin{aligned}
&\mathbb{E}\Big(\Big|e^{-cA} - e^{-cZ}\Big| \;\Big|\; Y_T > 0\Big) \\
&= \mathbb{E}\Big(\frac{Y_T - 1}{Y_T}\mathbb{E}\Big(\Big|e^{-cA} - e^{-cZ}\Big| \;\Big|\; Y_T, J_T < Y_T\Big) + \frac{1}{Y_T}\mathbb{E}\Big(\Big|e^{-cA} - e^{-cZ}\Big| \;\Big|\; Y_T, J_T = Y_T\Big) \;\Big|\; Y_T > 0\Big) \\
&\leq \mathbb{E}\Big(\mathbb{E}\Big(\Big|e^{-cA} - e^{-cZ}\Big| \;\Big|\; Y_T, J_T < Y_T\Big) + \frac{1}{Y_T} \;\Big|\; Y_T > 0\Big).
\end{aligned} \tag{17}$$

By Remark 3.17, the $\mathrm{Exp}(\lambda)$ distribution stochastically dominates the distribution corresponding to F_*. Thus we have $F_\infty(A) \le F_*(A)$. Moreover, given $J_T < Y_T$, we have $Z = F_\infty^{-1}(F_*(A))$. Since F_∞^{-1} is increasing, this implies that, given $J_T < Y_T$, we have $A \le Z$ almost surely. Thus the first summand in the outer expectation on the right-hand side of (17) is equal to

$$\mathbb{E}\left(e^{-cA} - e^{-cZ} \,\middle|\, Y_T, J_T < Y_T\right) = \mathbb{E}(e^{-cA}|Y_T, J_T < Y_T) - \mathbb{E}(e^{-cZ}|Y_T, J_T < Y_T). \tag{18}$$

Since, given $Y_T = y_T > 0$ and $J_T < y_T$, the random variable Z is $\mathrm{Exp}(\lambda)$ distributed, we have

$$\mathbb{E}(e^{-cZ}|J_T < Y_T) = \frac{\lambda}{\lambda + c}.$$

For the first expectation on the right-hand side of (18), we compute

$$\begin{aligned}\mathbb{E}(e^{-cA}|Y_T, J_T < Y_T) &= \frac{1}{1 - e^{-(\lambda-\mu)T}} \int_0^T (\lambda e^{-(\lambda+c)t} - \mu e^{-(\lambda-\mu)T} e^{-(\mu+c)t}) dt \\ &= \frac{1}{1 - e^{-(\lambda-\mu)T}} \left(\frac{\lambda}{\lambda + c}(1 - e^{-(\lambda+c)T}) - \frac{\mu}{\mu + c} e^{-(\lambda-\mu)T}(1 - e^{-(\mu+c)T}) \right) \\ &\le \frac{\lambda}{\lambda + c} \frac{e^{(\lambda-\mu)T}}{e^{(\lambda-\mu)T} - 1},\end{aligned}$$

where we assume $\lambda > \mu$. Thus for $\lambda > \mu$, the right-hand side of (18) is smaller than or equal to

$$\frac{\lambda}{\lambda + c}\left(\frac{e^{(\lambda-\mu)T}}{e^{(\lambda-\mu)T} - 1} - 1 \right) \le \frac{\lambda}{c + \lambda} \frac{1}{e^{(\lambda-\mu)T} - 1}. \tag{19}$$

For the second summand in the outer expectation on the right-hand side of (17), we use the upper bound from Proposition 3.3.

In order to prove Theorem 3.12, it remains to compute the age distribution of the last individual given $Y_T = y_T$. We do this analogously to the procedure for the first $y_T - 1$ individuals.

Recall from (12) that we have to compute $\mathcal{L}(\tilde{H}_{\tilde{W}_2} \mid \max_{x\ge 0} \tilde{H}_x = T)$. Note that $\tilde{H}_{\tilde{W}_2}$ has a density $\tilde{f}$ with respect to the measure $Leb_{[0,T)} + \delta_T$ given by

$$\tilde{f}(t) = \begin{cases} \lambda e^{-\lambda t} & \text{if } 0 \le t < T \\ e^{-\lambda T} & \text{if } t = T. \end{cases}$$

Thus by Bayes' Theorem, the age distribution of the last individual given $Y_T = y_T$ has a density f^* with respect to $Leb_{[0,T)} + \delta_T$ satisfying

$$f^*(t) \propto \tilde{f}(t) \mathbb{P}\left(\max_{x\ge 0} \tilde{H}_x = T \,\middle|\, \tilde{H}_{\tilde{W}_2} = t \right) = \begin{cases} \lambda e^{-\lambda t} \mathbb{P}\left(\max_{x\ge 0} \tilde{H}_x = T \,\middle|\, \tilde{H}_{\tilde{W}_2} = t \right) & \text{if } 0 \le t < T \\ e^{-\lambda T} & \text{if } t = T. \end{cases}$$

Firstly, we consider the case where $\lambda \ne \mu$. By (14), we have that

$$\mathbb{P}\left(\max_{x\ge 0} \tilde{H}_x = T \,\middle|\, \tilde{H}_{\tilde{W}_2} = t \right) = 1 - \mathbb{P}\left(\max_{x\ge 0} \tilde{H}_x < T \,\middle|\, \tilde{H}_{\tilde{W}_2} = t \right) = 1 - \frac{\lambda e^{(\lambda-\mu)(T-t)} - \mu}{\lambda e^{(\lambda-\mu)T} - \mu} e^{(\lambda-\mu)t}$$

for $t \in [0,T)$, which implies

$$f^*(t) \propto \begin{cases} \lambda e^{-\lambda t} \frac{\mu e^{(\lambda-\mu)t} - \mu}{\lambda e^{(\lambda-\mu)T} - \mu} & \text{if } 0 \leq t < T \\ e^{-\lambda T} & \text{if } t = T. \end{cases}$$

We can compute the normalizing constant of f^* as

$$\begin{aligned} e^{-\lambda T} + \frac{\lambda\mu}{\lambda e^{(\lambda-\mu)T} - \mu} \int_0^T e^{-\lambda t}(e^{(\lambda-\mu)t} - 1)dt &= e^{-\lambda T} + \frac{\lambda\mu}{\lambda e^{(\lambda-\mu)T} - \mu} \int_0^T (e^{-\mu t} - e^{-\lambda t})dt \\ &= e^{-\lambda T} + \frac{\lambda(1 - e^{-\mu T}) - \mu(1 - e^{-\lambda T})}{\lambda e^{(\lambda-\mu)T} - \mu} \\ &= \frac{\lambda - \mu}{\lambda e^{(\lambda-\mu)T} - \mu}. \end{aligned} \tag{20}$$

3.18 Remark

By Bayes' Theorem, the normalizing constant of the density f^* is just the probability that a linear birth and death process with birth rate μ and death rate λ survives up to time T. Thus we could also have used the extinction probability from Subsection 3.1 in order to obtain the right-hand side of (20).

We may conclude that f^* is given by

$$f^*(t) = \begin{cases} \lambda\mu \frac{e^{-\mu t} - e^{-\lambda t}}{\lambda - \mu} & \text{if } 0 \leq t < T \\ \frac{\lambda e^{-\mu T} - \mu e^{-\lambda T}}{\lambda - \mu} & \text{if } t = T. \end{cases}$$

By integration, we obtain that the cumulative distribution function F^* for the age of the last individual given $Y_T = y_T$ takes the form

$$F^*(t) = \frac{\lambda(1 - e^{-\mu t}) - \mu(1 - e^{-\lambda t})}{\lambda - \mu} \mathbb{1}_{\{t<T\}} + \mathbb{1}_{\{t=T\}} \tag{21}$$

for $t \in [0,T]$.

For $\lambda = \mu > 0$, we proceed analogously: By (15), we have that

$$\mathbb{P}\left(\max_{x\geq 0} \tilde{H}_x = T \,\middle|\, \tilde{H}_{\tilde{W}_2} = t\right) = 1 - \mathbb{P}\left(\max_{x\geq 0} \tilde{H}_x < T \,\middle|\, \tilde{H}_{\tilde{W}_2} = t\right) = 1 - \frac{1 + \lambda(T-t)}{1 + \lambda T} = \frac{\lambda t}{1 + \lambda T}$$

for $t \in [0,T)$, which leads to

$$f^*(t) \propto \begin{cases} \lambda e^{-\lambda t} \frac{\lambda t}{1+\lambda T} & \text{if } 0 \leq t < T \\ e^{-\lambda T} & \text{if } t = T. \end{cases}$$

We compute the normalizing constant of f^* as

$$e^{-\lambda T} + \int_0^T e^{-\lambda t} \frac{\lambda t}{1 + \lambda T} dt = e^{-\lambda T} + \frac{1}{1 + \lambda T} - e^{-\lambda T} = \frac{1}{1 + \lambda T}.$$

This yields that f^* is given by

$$f^*(t) = \begin{cases} \lambda^2 t e^{-\lambda t} & \text{if } 0 \leq t < T \\ (1+\lambda T)e^{-\lambda T} & \text{if } t = T. \end{cases}$$

By integration, we obtain that the cumulative distribution function F^* for the age of the last individual given $Y_T = y_T$ takes the form

$$F^*(t) = (1 - e^{-\lambda t}(1+\lambda t)\mathbb{1}_{\{t<T\}} + \mathbb{1}_{\{t=T\}}$$

for $t \in [0, T]$.

By multiplying the probability that given $Y_T = y_T$ the randomly picked individual is the last one, we obtain the second summands in Theorem 3.12. $\square$

3.4 Upper bound for the time since the last event

Let $0 = T_1 < T_2 < \dots$ be the event times of our linear birth and death process. Since $B_T + D_T$ is the number of events up to time T, the random variable $T - T_{B_T+D_T}$ describes the time passed since the last event as seen from time T. Note that, given the population size $Y_T = y_T$, the distribution of the difference $T_{B_T+D_T+1} - T_{B_T+D_T}$ between the previous and the next event time at time T is typically larger than the $\text{Exp}(y_T(\lambda+\mu))$ distribution due to the inspection paradox. Therefore, we do not bound $T - T_{B_T+D_T}$ from above by $T_{B_T+D_T+1} - T_{B_T+D_T}$. Instead we use that, given $Y_T = y_T \geq 2$, the time since the last event is bounded from above by the minimum of the ages of the first $y_T - 1$ individuals, where the individuals are ordered as in the previous subsection. We know that these $y_T - 1$ ages are are i.i.d. and that their distribution is stochastically dominated by the $\text{Exp}(\lambda)$ distribution by Remark 3.17. Since the minimum of $y_T - 1$ independent $\text{Exp}(\lambda)$ distributed random variables, is $\text{Exp}((y_T-1)\lambda)$ distributed, we obtain the following theorem.

3.19 Theorem
Given $Y_T = y_T$, the distribution of $T - T_{B_T+D_T}$ is stochastically dominated by the distribution with cumulative distribution function

$$G(t) = \mathbb{1}_{\{y_T>1\}}(1 - e^{-(y_T-1)\lambda t}) + \mathbb{1}_{\{y_T\leq 1\}}\mathbb{1}_{\{t\geq T\}}.$$

4 Degree distribution at finite time

In this section, we deduce the degree distribution at finite time T.

Firstly, we consider the case where the nodes do not die, i.e. $\mu = 0$. Let the nodes be ordered by their birth times. Let S_i be the social index of node i and $A_i(T)$ its age at time T. For convenience, we define $A_2(T) = 0$ if $Y_T = 1$. Furthermore, given $Y_T = y_T$, let the random variable J_T be uniformly distributed on $\{1, \dots, y_T\}$ and independent of the ages, the social indices and the rest of the path $(Y_t)_{0\leq t<T}$. We interpret J_T as the index of a node that is randomly chosen at time T among all living nodes.

4.1 The number of outgoing edges in the pure birth case

Firstly, we condition on the population size y_T, the randomly picked node j_T, its social index s_{j_T} and its age a_{j_T} at time T. Furthermore, we condition on the age a_2 of the second node if $y_T > 1$ and set $a_2 = 0$ otherwise. If $y_T > 1$, the number of living edges that j_T creates during its lifetime can be described by a birth and death process with constant birth and linear death rate started in zero at time $T - a_{\max(j_T,2)}$ since no edges are created as long as there is only one node. The birth rate is αs_{j_T} and the death rate is the product of β and the number of (living) edges that j_T has created. Such a birth and death process is examined in Chapter XVII of [Fel67] and can be interpreted as a queueing system with constant arrival rate and infinitely many servers. We denote the state probabilities at time t of such a process started at a general state i with constant birth rate $\tilde{\lambda}$ and linear death rate $n\tilde{\mu}$ by $P_n(t)$, where n is the current state of the process. This process is described by the following Kolmogorov forward differential equations:

$$\frac{dP_n(t)}{dt} = \tilde{\lambda} P_{n-1}(t) - (\tilde{\lambda} + n\tilde{\mu})P_n(t) + \tilde{\mu}(n+1)P_{n+1}(t) \text{ for } n \geq 1 \tag{22}$$

$$\frac{P_0(t)}{dt} = \tilde{\mu}P_1(t) - \tilde{\lambda}P_0(t). \tag{23}$$

Thus we have for the generating function $P(s,t) = \sum\limits_{n=0}^{\infty} P_n(t)s^n$ for all s with $|s| < 1$:

$$\begin{aligned}\frac{\partial P(s,t)}{\partial t} &= \frac{\partial \sum\limits_{n=0}^{\infty} P_n(t)s^n}{\partial t} = \sum_{n=0}^{\infty} \frac{dP_n(t)}{dt} s^n \\ &= \sum_{n=1}^{\infty} s^n \Big(\tilde{\lambda} P_{n-1}(t) - (\tilde{\lambda} + n\tilde{\mu})P_n(t) + \tilde{\mu}(n+1)P_{n+1}(t) \Big) + \tilde{\mu}P_1(t) - \tilde{\lambda}P_0(t) \\ &= (1-s)\Big(-\tilde{\lambda}P(s,t) + \tilde{\mu}\frac{\partial P(s,t)}{\partial s} \Big), \end{aligned} \tag{24}$$

where the second line follows from the Kolmogorov forward differential equations (22) and (23). The solution of the partial differential equation (24) is given by (cf. [Fel67], page 481)

$$P(s,t) = \exp\Big(-\tilde{\lambda}(1-s)\frac{(1-e^{-\tilde{\mu}t})}{\tilde{\mu}} \Big)(1-(1-s)e^{-\tilde{\mu}t})^i. \tag{25}$$

The function $P(\,\cdot\,, t)$ is the product of the generating function of the $\mathrm{Po}(\frac{\tilde{\lambda}}{\tilde{\mu}}(1-e^{-\tilde{\mu}t}))$ and the $\mathrm{Bin}(i, e^{-\tilde{\mu}t})$ distribution and hence the generating function of their convolution. In particular, we obtain for $i = 0$ the generating function of the Poisson distribution with parameter

$$\frac{\tilde{\lambda}(1-e^{-\tilde{\mu}t})}{\tilde{\mu}}.$$

This result can be applied to the edge process we are interested in: Conditioned on the population size y_T, the chosen node j_T, its social index s_{j_T} and age a_{j_T}, the distribution of the number of edges created by j_T is the following Poisson distribution:

$$\mathrm{Po}\Big(\frac{\alpha s_{j_T}}{\beta}(1 - e^{-\beta a_{\max(j_T,2)}}) \Big),$$

where the Po(0) distribution is the Dirac measure in zero. Note that we obtain the Dirac measure if only one individual is alive at time T, i.e. $y_T = 1$, since we set $a_2 = 0$ in this case.

Consequently, the corresponding unconditional distribution is the following mixed Poisson distribution:

$$\mathrm{MixPo}\bigg(\frac{\alpha S_{J_T}}{\beta}(1 - e^{-\beta A_{\max(J_T,2)}(T)})\bigg).$$

4.2 The number of incoming edges in the pure birth case

Recall that the nodes are ordered by their birth times. Firstly, we condition on $(Y_t)_{0\le t\le T} = (y_t)_{0\le t\le T}$, the social indices $(S_k)_{k\in\mathbb{N}} = (s_k)_{k\in\mathbb{N}}$ and $J_T = j_T$. Let $T = a_1 > \ldots > a_{y_T}$ denote the corresponding ages of the individuals.

Consider a fixed node $i \in \{1, \ldots, y_T\} \setminus \{j_T\}$ and some time interval of the form $[T - a_l, T - a_{l+1})$ for $l \ge i \vee 2$. We only consider edges that are created by the fixed node i. Then the number of edges that connect i to j_T and that survive until the end of this interval, i.e. until the birth time $T - a_{l+1}$ of node $l + 1$, can be described by a birth and death process again. The population size is constant and equal to l in the interval $[T - a_l, T - a_{l+1})$. Thus the birth rate is also constant and equal to $\alpha s_i \frac{1}{l-1}$ because node i creates edges at rate αs_i and $\frac{1}{l-1}$ is the probability that an edge that is created in the interval $[T - a_l, T - a_{l+1})$ is connected to j_T. The death rate is linear again with factor β. As in the previous subsection, we obtain a Poisson distribution, i.e. the number of edges that are created by i in $[T - a_l, T - a_{l+1})$, connect i to j_T and survive until $T - a_{l+1}$ follows the distribution

$$\mathrm{Po}\bigg(\frac{\alpha s_i}{(l-1)\beta}(1 - e^{-\beta(a_l - a_{l+1})})\bigg).$$

Now the distribution of the number of edges that i creates in $[T - a_l, T - a_{l+2})$, connect i with j_T and survive until $T - a_{l+2}$ can be deduced from (25). In order to do this, let Z be the number of edges that i creates in $[T - a_l, T - a_{l+1})$, connect i with j_T and survive until $T - a_{l+1}$. We have already shown that Z is $\mathrm{Po}(\frac{\alpha s_i}{(l-1)\beta}(1 - e^{-\beta(a_l - a_{l+1})}))$ distributed. First, we condition on $Z = z$. Then the number of edges that i creates in $[T - a_l, T - a_{l+2})$, connect i with j_T and survive until $T - a_{l+2}$ is described by a birth and death process with constant birth rate $\frac{\alpha s_i}{l}$, linear death rate with factor β and initial state z. For such a process, the probability generating function (25) was stated in the previous subsection. Since (25) is the product of the probability generating function of a Poisson distribution and of the probability generating function of a binomial distribution, we obtain the sum of a $\mathrm{Po}(\frac{\alpha s_i}{l\beta}(1 - e^{-\beta(a_{l+1} - a_{l+2})}))$ distributed and a $\mathrm{Bin}(z, e^{-\beta(a_{l+1} - a_{l+2})})$ distributed random variable for time $T - a_{l+2}$, where these two random variables are independent. The second one is a Poisson mixture of binomial distributions if we do not condition on $Z = z$. This mixture is the Poisson distribution with parameter $e^{-\beta(a_{l+1} - a_{l+2})}\frac{\alpha s_i}{(l-1)\beta}(1 - e^{-\beta(a_l - a_{l+1})})$. Consequently, it follows that

$$\mathrm{Po}\bigg(e^{-\beta(a_{l+1} - a_{l+2})}\frac{\alpha s_i}{(l-1)\beta}(1 - e^{-\beta(a_l - a_{l+1})}) + \frac{\alpha s_i}{l\beta}(1 - e^{-\beta(a_{l+1} - a_{l+2})})\bigg).$$

is the desired distribution.

We obtain inductively by mixing and independence that

$$\mathrm{Po}\Big(\sum_{l=i\vee j_T}^{y_T-1}\Big(\prod_{k=1}^{y_T-l-1} e^{-\beta(a_{l+k}-a_{l+k+1})}\Big)\frac{\alpha s_i}{(l-1)\beta}(1-e^{-\beta(a_l-a_{l+1})})\Big)$$

is the distribution of the number of edges that connect i to j_T surviving until $T-a_{y_T}$.

Repeating the same argument as above, it follows that

$$\mathrm{Po}\Big(\frac{\alpha s_i}{(y_T-1)\beta}(1-e^{-\beta a_{y_T}})+e^{-\beta a_{y_T}}\sum_{l=i\vee j_T}^{y_T-1}\Big(\prod_{k=1}^{y_T-l-1} e^{-\beta(a_{l+k}-a_{l+k+1})}\Big)\frac{\alpha s_i}{(l-1)\beta}(1-e^{-\beta(a_l-a_{l+1})})\Big)$$

is the distribution of the number of edges between i and j_T being alive at time T.

By independence of the edge processes of edges generated by the various nodes i, we obtain the following distribution for the total number of incoming edges in node j_T:

$$\begin{aligned}&\mathrm{Po}\Big(\sum_{\substack{i=1\\ i\neq j_T}}^{y_T}\frac{\alpha s_i}{(y_T-1)\beta}(1-e^{-\beta a_{y_T}})+\sum_{\substack{i=1\\ i\neq j_T}}^{y_T} e^{-\beta a_{y_T}}\sum_{l=i\vee j_T}^{y_T-1}\Big(\prod_{k=1}^{y_t-l-1} e^{-\beta(a_{l+k}-a_{l+k+1})}\Big)\frac{\alpha s_i}{(l-1)\beta}(1-e^{-\beta(a_l-a_{l+1})})\Big)\\ &=\mathrm{Po}\Big(\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq j_T}}^{y_T}\frac{s_i}{y_T-1}(1-e^{-\beta a_{y_T}})+\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq j_T}}^{y_T}\sum_{l=i\vee j_T}^{y_T-1}\frac{s_i}{l-1}(e^{-\beta a_{l+1}}-e^{-\beta a_l})\Big).\end{aligned}$$

Thus without conditioning, we obtain a mixed Poisson distribution:

$$\mathrm{MixPo}\Big(\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{Y_T}\frac{S_i}{Y_T-1}(1-e^{-\beta A_{Y_T}(T)})+\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{Y_T}\sum_{l=i\vee J_T}^{Y_T-1}\frac{S_i}{l-1}(e^{-\beta A_{l+1}(T)}-e^{-\beta A_l(T)})\Big).$$

4.3 Degree distribution in the pure birth case

By the independence of the numbers of incoming and outgoing edges, the previous considerations imply that the degree distribution of a randomly chosen node is a mixed Poisson distribution. The random parameter of this mixed Poisson distribution is the sum of the random parameters of the two mixed Poisson distributions that were deduced in the previous two subsections. Thus we obtain the $\mathrm{MixPo}(\Lambda_T)$ distribution, where

$$\begin{aligned}\Lambda_T=&\frac{\alpha S_{J_T}}{\beta}(1-e^{-\beta A_{\max(J_T,2)}(T)})+\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{Y_T}\frac{S_i}{Y_T-1}(1-e^{-\beta A_{Y_T}(T)})\\ &+\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{Y_T}\sum_{l=i\vee J_T}^{Y_T-1}\frac{S_i}{l-1}(e^{-\beta A_{l+1}(T)}-e^{-\beta A_l(T)}).\end{aligned}\tag{26}$$

4.4 Degree distribution in the general case

If the nodes can die, the degree distribution can be determined similarly to the pure birth case.

Let $0 = T_1^+ < \ldots < T_{B_T}^+$ be the birth times. Recall that $0 = T_1 < \ldots < T_{B_T+D_T}$ are the event times up to T. Moreover, let T_i^- denote the death time of node i for each i, where the nodes are still numbered according to their birth times. Finally, given the subset of $\{1, \ldots, B_T\}$ that contains the indices of all living nodes at time T, let J_T be uniformly distributed on this set and independent of all other events as before. Note that S_{J_T} is (stochastically) independent of $((Y_t)_{0\le t\le T}, J_T)$.

The crucial differences compared to the procedure for the pure birth case are:

- We have to condition on $Y_T > 0$ since there has to exist a node at time T that can be picked randomly.
- We have to consider $[T_l, T_{l+1})$ instead of $[T - A_l(T), T - A_{l+1}(T))$ as time intervals in which the parameters of the process of incoming edges are constant.
- We do not sum over all nodes i, but over all that are still alive at time T.
- In general, the population size does not increase almost surely by one per time interval. Thus the population size is not necessarily equal to l in the time interval $[T_l, T_{l+1})$.

Since deaths of other nodes reduce the number of outgoing edges, we cannot derive the distribution of the number of outgoing edges at time T in the same way as for the pure birth process. However, we can proceed in a similar way as for the incoming nodes in the pure birth case. Therefore we do not derive the distribution of the numbers of outgoing and incoming edges separately here.

We condition on $(Y_t)_{0\le t\le T} = (y_t)_{0\le t\le T}$, the social indices $(S_k)_{k\in\mathbb{N}} = (s_k)_{k\in\mathbb{N}}$ and $J_T = j_T$. Let b_T and d_T denote the corresponding number of births and deaths up to time T, respectively. Furthermore, let $0 = t_1 < t_2 < \ldots < t_{b_T+d_T}$ and $0 = t_1^+ < t_2^+ < \ldots < t_{b_T}^+$ be the corresponding event times and birth times up to time T, respectively. We consider a fixed node $i \neq j_T$ that is alive at time T (provided there are any) and some time interval of the form $[t_l, t_{l+1})$ for $t_l \ge t_i^+ \vee t_{j_T}^+$ and $t_{l+1} \le T$. Note that the nodes i and j_T create edges with rate αs_i and αs_{j_T}, respectively, and that the probability that an edge that is created by i is connected to j_T is $\frac{1}{y_{t_l}-1}$ and equal to the probability that an edge that is created by j_T is connected to i. Thus the number of edges that connect i to j_T and survive until time t_{l+1} can be described by a birth and death process with constant birth rate $\alpha(s_i + s_{j_T})\frac{1}{y_{t_l}-1}$ and linear death rate with factor β like before. Analogously to Subsection 4.2, we obtain that the number of edges that are created in $[t_l, t_{l+1})$, connect i to j_T and survive until t_{l+1} has the following Poisson distribution:

$$\mathrm{Po}\biggl(\frac{\alpha(s_i + s_{j_T})}{(y_{t_l} - 1)\beta}(1 - e^{-\beta(t_{l+1}-t_l)})\mathbb{1}_{\{y_{t_l}>1\}}\biggr).$$

Let the function $r = r_{(y_t)_{0\le t\le T}} : \{1, \ldots, b_T\} \to \{1, \ldots, b_T + d_T\}$ be defined such that $t_j^+ = t_{r(j)}$ for all $j \in \{1, \ldots, b_T\}$, i.e. r maps birth number to event number. Note that the sum $b_T + d_T$ of the number of births and the number of deaths up to time T is the number of events up to time T.

Applying the same iterative procedure as in Subsection 4.2, we can see that

$$\begin{aligned}&\mathrm{Po}\bigg(\frac{\alpha(s_i+s_{j_T})}{(y_T-1)\beta}(1-e^{-\beta(T-t_{b_T+d_T})})\mathbb{1}_{\{y_T>1\}}\\&\quad+e^{-\beta(T-t_{b_T+d_T})}\sum_{l=r(i)\vee r(j_T)}^{b_T+d_T-1}\bigg(\prod_{k=1}^{b_T+d_T-l-1}e^{-\beta(t_{l+k+1}-t_{l+k})}\frac{\alpha(s_i+s_{j_T})}{(y_{t_l}-1)\beta}(1-e^{-\beta(t_{l+1}-t_l)})\bigg)\mathbb{1}_{\{y_{t_l}>1\}}\bigg)\end{aligned}$$

is the distribution of the number of edges between i and j_T being alive at time T.

Let the death times be denoted by $t_1^-, t_2^-, \dots, t_{d_T}^-$. Then by independence of the processes of edges generated by the various nodes i, we obtain the following conditional degree distribution:

$$\begin{aligned}&\mathrm{Po}\bigg(\sum_{\substack{i=1\\i\neq j_T}}^{b_T}\frac{\alpha(s_i+s_{j_T})}{(y_T-1)\beta}\mathbb{1}_{\{t_i^->T\}}(1-e^{-\beta(T-t_{b_T+d_T})})\mathbb{1}_{\{y_T>1\}}+\sum_{\substack{i=1\\i\neq j_T}}^{b_T}\mathbb{1}_{\{t_i^->T\}}e^{-\beta(T-t_{b_T+d_T})}\\&\quad\cdot\sum_{l=r(i)\vee r(j_T)}^{b_T+d_T-1}\bigg(\bigg(\prod_{k=1}^{b_T+d_T-l-1}e^{-\beta(t_{l+k+1}-t_{l+k})}\bigg)\frac{\alpha(s_i+s_{j_T})}{(y_{t_l}-1)\beta}(1-e^{-\beta(t_{l+1}-t_l)})\bigg)\mathbb{1}_{\{y_{t_l}>1\}}\bigg)\\&=\mathrm{Po}\bigg(\frac{\alpha}{\beta}\sum_{\substack{i=1\\i\neq j_T}}^{b_T}\frac{s_i+s_{j_T}}{(y_T-1)}\mathbb{1}_{\{t_i^->T\}}(1-e^{-\beta(T-t_{b_T+d_T})})\mathbb{1}_{\{y_T>1\}}\\&\quad+\frac{\alpha}{\beta}\sum_{\substack{i=1\\i\neq j_T}}^{b_T}\mathbb{1}_{\{t_i^->T\}}\sum_{l=r(i)\vee r(j_T)}^{b_T+d_T-1}\frac{s_i+s_{j_T}}{(y_{t_l}-1)}(e^{-\beta(T-t_{l+1})}-e^{-\beta(T-t_l)})\mathbb{1}_{\{y_{t_l}>1\}}\bigg).\end{aligned}$$

Thus the degree distribution is $\mathrm{MixPo}(\Lambda_T^*)$, where $\mathcal{L}(\Lambda_T^*)=\mathcal{L}(\Lambda_T|Y_T>0)$ and

$$\begin{aligned}\Lambda_T:=&\frac{\alpha}{\beta}\sum_{\substack{i=1\\i\neq J_T}}^{B_T}\frac{S_i+S_{J_T}}{(Y_T-1)}\mathbb{1}_{\{T_i^->T\}}(1-e^{-\beta(T-T_{B_T+D_T})})\mathbb{1}_{\{Y_T>1\}}\\&+\frac{\alpha}{\beta}\sum_{\substack{i=1\\i\neq J_T}}^{B_T}\mathbb{1}_{\{T_i^->T\}}\sum_{l=r(i)\vee r(J_T)}^{B_T+D_T-1}\frac{S_i+S_{J_T}}{(Y_{T_l}-1)}(e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)})\mathbb{1}_{\{Y_{T_l}>1\}}.\end{aligned}\tag{27}$$

5 Bounds on the total variation distance between the finite time and asymptotic degree distribution

Since in the pure birth case we obtain a much better bound with considerably less work, we treat the cases $\mu=0$ and $\mu>0$ separately.

5.1 The pure birth case

Firstly, we consider the pure birth case, i.e. $\mu=0$, again. Fix $T>0$ and let the random variable Λ_T be defined by (26). Furthermore, let S_{J_∞} and A_{J_∞} be independent random variables such that $S_{J_\infty}(\omega)=S_{J_T}(\omega)$ and $A_{J_\infty}(\omega):=F_\infty^{-1}(F_T(A_{J_T}(T,\omega)))$ for all $\omega\in\Omega$, where F_T and F_∞ are the cumulative distribution functions of $A_{J_T}(T)$ and the $\mathrm{Exp}(\lambda)$ distribution, respectively. Note that we obtain from Proposition 3.9 that $\mathcal{L}(A_{J_T}) \to \text{Exp}(\lambda)$ weakly as $T \to \infty$. The random variable A_{J_∞} is $\text{Exp}(\lambda)$ distributed since F_T is continuous and hence $F_T(A_{J_T}(T))$ is uniformly distributed on $[0,1]$. Moreover, since the exponential distribution dominates the truncated exponential distribution, we have $F_\infty(A_{J_T}(T,\omega)) \leq F_T(A_{J_T}(T,\omega))$ for all $\omega \in \Omega$. Since F_∞^{-1} is increasing, it follows that $A_{J_T}(T,\omega) \leq A_{J_\infty}(\omega)$ for all $\omega \in \Omega$.

Let S be a generic random variable that is distributed according to the distribution of the social indices and let

$$\mathrm{M} := \frac{\alpha S_{J_\infty}}{\beta}(1 - e^{-\beta A_{J_\infty}}) + \frac{\alpha \mathbb{E}(S)}{\beta}(1 - e^{-\beta A_{J_\infty}}). \tag{28}$$

We know from Subsection 4.3 that $\text{MixPo}(\Lambda_T)$ is the degree distribution at time T in the pure birth case. Theorem 5.2 below implies that $\text{MixPo}(\Lambda_T)$ converges at rate of just a bit slower than $e^{-\frac{1}{2}\lambda T}$ to $\text{MixPo}(\mathrm{M})$ as $T \to \infty$, which is the asymptotic degree distribution already stated in Section 3.2 of [BL10]. Note that these theorems are much more powerful as they give an exact distance bound for every finite T.

5.1 Theorem
We have

$$d_{TV}(\text{MixPo}(\Lambda_T), \text{MixPo}(\mathrm{M})) \leq \frac{4\alpha}{\lambda}\sigma_S \mathbb{E}\left(\frac{1}{\sqrt{Y_T}}\right) + \frac{2\alpha}{\lambda}\mathbb{E}(S)\mathbb{E}\left(\frac{1}{Y_T}\right) + \frac{2\alpha}{\beta}\mathbb{E}(S)\frac{\lambda}{\beta+\lambda}\frac{1}{e^{\lambda T}-1},$$

where σ_S is the standard deviation of S.

The main idea of the proof is to make use of Theorem 2.1 to obtain $\mathbb{E}(|\Lambda_T - \mathrm{M}|)$ as an upper bound for the total variation distance on the left-hand side. In order to bound $\mathbb{E}(|\Lambda_T - \mathrm{M}|)$ further, we use that the expected value of the second summand of the right-hand side of (26) becomes small as $T \to \infty$ since the age $A_{Y_T}(T)$ of the youngest individual at time T converges quickly to 0, and compare the other summands of the right-hand sides of (26) and (28).

For the comparison of the last summands in (26) and (28), respectively, we note that the average of the social indices becomes close to $\mathbb{E}(S)$ by the Law of Large Numbers. Then we use again that the age $A_{Y_T}(T)$ of the youngest individual at time T converges quickly to 0 and that $A_{J_T}(T)$ converges quickly to A_{J_∞} Proposition 3.9.

Finally, the expected absolute value of the difference between the first summand of (26) and the first summand of (28) again becomes small since $A_{J_T}(T)$ converges quickly to A_{J_∞}.

We prove Theorem 5.1 in detail after the next theorem, which gives us the exponential rate for the convergence of the degree distribution. In order to reveal this rate, we further estimate $\mathbb{E}(\frac{1}{Y_T})$ and $\mathbb{E}(\frac{1}{\sqrt{Y_T}})$ on the right-hand side in Theorem 5.1 by using Proposition 3.3 and Corollary 3.4.

5.2 Theorem
For $T \geq \frac{\log(2)}{\lambda}$, we have

$$d_{TV}(\text{MixPo}(\Lambda_T), \text{MixPo}(\mathrm{M})) \leq \frac{\sqrt{32}\alpha}{\sqrt{\lambda}}\sigma_S\sqrt{T}e^{-\frac{1}{2}\lambda T} + 4\alpha\mathbb{E}(S)\left(T + \frac{\lambda}{\beta(\beta+\lambda)}\right)e^{-\lambda T},$$

where σ_S is the standard deviation of S.

Proof: The inequality follows directly from Theorem 5.1, Proposition 3.3 and Corollary 3.4. ☐

5.3 Remark
For the original model of Britton and Lindholm, which allows loops, we can adapt the proof of Theorem 5.1 in such a way that the upper bounds in Theorem 5.1 and Theorem 5.2 remain exactly the same.

Proof of Theorem 5.1

From Theorem 2.1 follows

$$d_{TV}(\mathrm{MixPo}(\Lambda_T), \mathrm{MixPo}(\mathrm{M})) \leq \mathbb{E}(|\mathrm{M} - \Lambda_T|),$$

and $\mathbb{E}(|\mathrm{M} - \Lambda_T|)$ is smaller than or equal to

$$\begin{aligned}
&\mathbb{E}\bigg|\frac{\alpha}{\beta}\bigg(S_{J_\infty}(1-e^{-\beta A_{J_\infty}}) + \mathbb{E}(S)(1-e^{-\beta A_{J_\infty}}) - S_{J_T}(1-e^{-\beta A_{\max(J_T,2)}(T)}) \\
&\qquad - \sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \frac{S_i}{Y_T-1}(1-e^{-\beta A_{Y_T}(T)}) - \sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \sum_{l=i\vee J_T}^{Y_T-1} \frac{S_i}{l-1}(e^{-\beta A_{l+1}(T)} - e^{-\beta A_l(T)})\bigg)\bigg| \\
&\leq \frac{\alpha}{\beta}\mathbb{E}\bigg|\sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \frac{S_i}{Y_T-1}(1-e^{-\beta A_{Y_T}(T)})\bigg| \\
&\quad + \frac{\alpha}{\beta}\mathbb{E}\bigg|\sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \sum_{l=i}^{Y_T-1} \frac{(S_i - \mathbb{E}(S))}{l-1}\mathbb{1}_{\{J_T\leq l\}}(e^{-\beta A_{l+1}(T)} - e^{-\beta A_l(T)})\bigg| \\
&\quad + \frac{\alpha}{\beta}\mathbb{E}\bigg|\mathbb{E}(S)(1-e^{-\beta A_{J_\infty}}) - \sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \sum_{l=i\vee J_T}^{Y_T-1} \frac{\mathbb{E}(S)}{l-1}(e^{-\beta A_{l+1}(T)} - e^{-\beta A_l(T)})\bigg| \\
&\quad + \frac{\alpha}{\beta}\mathbb{E}\bigg|S_{J_\infty}(1-e^{-\beta A_{J_\infty}}) - S_{J_T}(1-e^{-\beta A_{\max(J_T,2)}(T)})\bigg|, \qquad (29)
\end{aligned}$$

where we use the convention $\frac{0}{0} := 0$.

For the first line of the right-hand side, we have

$$\begin{aligned}
\mathbb{E}\bigg|\sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \frac{S_i}{Y_T-1}(1-e^{-\beta A_{Y_T}(T)})\bigg| &= \mathbb{E}\bigg(\mathbb{E}\bigg(\sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \frac{S_i}{Y_T-1}(1-e^{-\beta A_{Y_T}(T)})\bigg| Y_T, J_T\bigg)\bigg) \\
&= \mathbb{E}\bigg(\sum_{\substack{i=1 \\ i\neq J_T}}^{Y_T} \frac{\mathbb{E}(S_i)}{Y_T-1}(1-\mathbb{E}(e^{-\beta A_{Y_T}(T)}|Y_T))\bigg), \qquad (30)
\end{aligned}$$

where the last equality holds since, given Y_T, the age $A_{Y_T}(T)$ and J_T are independent.

Given $Y_T = y_T > 1$, the age $A_{y_T}(T)$ is the minimum of $y_T - 1$ i.i.d. truncated exponentially distributed random variables by Proposition 3.9. Since the minimum of $y_T - 1$ independent $\mathrm{Exp}(\lambda)$ distributed random variables is $\mathrm{Exp}((y_T-1)\lambda)$ distributed, the distribution of $A_{y_T}(T)$ is stochastically dominated by the $\mathrm{Exp}((y_T-1)\lambda)$ distribution. In general, we have for random variables $\hat{X}$ and $\hat{Y}$ with $\hat{X} \leq_{st} \hat{Y}$ that $f(\hat{X}) \leq_{st} f(\hat{Y})$ and consequently $\mathbb{E}(f(\hat{X})) \leq \mathbb{E}(f(\hat{Y}))$ for every increasing function f. Thus the right-hand side of (30) is smaller than or equal to

$$\mathbb{E}\bigg(\mathbb{E}(S)\frac{1}{\lambda}\frac{\beta}{Y_T-1+\frac{\beta}{\lambda}}\mathbb{1}_{\{Y_T>1\}}\bigg) \leq \frac{\beta}{\lambda}\mathbb{E}(S)\mathbb{E}\bigg(\frac{1}{Y_T-1}\mathbb{1}_{\{Y_T>1\}}\bigg).$$

Note that

$$\begin{aligned}\mathbb{E}\bigg(\frac{1}{Y_T-1}\mathbb{1}_{\{Y_T>1\}}\bigg) &= \sum_{n=2}^{\infty}\frac{p_n(T)}{n-1} = \frac{\lambda}{\lambda e^{\lambda T}}\sum_{n=2}^{\infty}\frac{1}{n-1}\bigg(\frac{\lambda e^{\lambda T}-\lambda}{\lambda e^{\lambda T}}\bigg)^{n-1} = \frac{\lambda}{\lambda e^{\lambda T}}\sum_{n=1}^{\infty}\frac{1}{n}\bigg(\frac{\lambda e^{\lambda T}-\lambda}{\lambda e^{\lambda T}}\bigg)^{n}\\ &\leq \frac{\lambda}{\lambda e^{\lambda T}-\lambda}\sum_{n=1}^{\infty}\frac{1}{n}\bigg(\frac{\lambda e^{\lambda T}-\lambda}{\lambda e^{\lambda T}}\bigg)^{n} = \sum_{n=1}^{\infty}\frac{p_n(T)}{n} = \mathbb{E}\bigg(\frac{1}{Y_T}\bigg),\end{aligned}$$

where $p_n(T)$ is the probability mass function from Subsection 3.1. Thus the right-hand side of (30) is smaller than or equal to

$$\frac{\beta}{\lambda}\mathbb{E}(S)\mathbb{E}\bigg(\frac{1}{Y_T}\bigg).$$

For the second line of the right-hand side of (29), we obtain

$$\begin{aligned}&\mathbb{E}\bigg|\sum_{\substack{i=1\\ i\neq J_T}}^{Y_T}\sum_{l=i}^{Y_T-1}\frac{(S_i-\mathbb{E}(S))}{l-1}\mathbb{1}_{\{J_T\leq l\}}(e^{-\beta A_{l+1}(T)}-e^{-\beta A_l(T)})\bigg|\\ &=\mathbb{E}\bigg|\sum_{l=2}^{Y_T-1}\frac{1}{l-1}\sum_{\substack{i=1\\ i\neq J_T}}^{l}(S_i-\mathbb{E}(S))\mathbb{1}_{\{J_T\leq l\}}(e^{-\beta A_{l+1}(T)}-e^{-\beta A_l(T)})\bigg|\\ &=\mathbb{E}\Bigg(\mathbb{E}\Bigg(\bigg|\sum_{l=J_T\vee 2}^{Y_T-1}\frac{1}{l-1}\sum_{\substack{i=1\\ i\neq J_T}}^{l}(S_i-\mathbb{E}(S))(e^{-\beta A_{l+1}(T)}-e^{-\beta A_l(T)})\bigg|\ \bigg|\ Y_T,J_T\Bigg)\Bigg)\\ &\leq\mathbb{E}\Bigg(\sum_{l=J_T\vee 2}^{Y_T-1}\mathbb{E}\bigg|\frac{1}{l-1}\sum_{i=2}^{l}(S_i-\mathbb{E}(S))\bigg|\mathbb{E}\Big(e^{-\beta A_{l+1}(T)}(1-e^{-\beta(A_l(T)-A_{l+1}(T))})|Y_T\Big)\Bigg)\\ &\leq\mathbb{E}\Bigg(\sum_{l=J_T\vee 2}^{Y_T-1}\frac{1}{l-1}\sqrt{\sum_{i=1}^{l}\mathbb{E}\Big((S_i-\mathbb{E}(S))^2\Big)}\mathbb{E}(1-e^{-\beta(A_l(T)-A_{l+1}(T))}|Y_T)\Bigg)\\ &\leq\mathbb{E}\Bigg(\sum_{l=J_T\vee 2}^{Y_T-1}\frac{1}{\sqrt{l-1}}\sigma_S\mathbb{E}(1-e^{-\beta(A_l(T)-A_{l+1}(T))}|Y_T)\Bigg),\end{aligned}\tag{31}$$

where the fourth line holds since the second sum in the third line is (stochastically) independent of J_T, the social indices are independent of all other random variables and, given Y_T, the index J_T is independent of the social indices and ages, and the second last line is obtained by applying $\mathbb{E}|Z-\mathbb{E}(Z)| \leq \sqrt{\mathrm{Var}(Z)}$ to $Z=\frac{1}{l-1}\sum_{i=2}^{l}S_i$.

Given Y_T and $A_{l+1}(T)$, the difference $A_l(T)-A_{l+1}(T)$ is the minimum of $l-1$ i.i.d. truncated exponentially distributed random variables by Proposition 3.9, Corollary 3.10 and Remark 3.11 (for

$l \geq 2$). Thus it is stochastically dominated by an $\mathrm{Exp}((l-1)\lambda)$ distributed random variable Z_l, where Z_l can be assumed to independent of $A_{l+1}(T)$. This implies for $a > 0$:

$$\begin{aligned}\mathbb{P}(A_l - A_{l+1} \leq a|Y_T) &= \int_0^T \mathbb{P}(A_l - A_{l+1} \leq a|Y_T, T - A_{l+1} = x)\mathbb{P}(T - A_{l+1} \in dx|Y_T) \\ &\geq \mathbb{P}(Z_l \leq a|Y_T).\end{aligned}$$

Thus, also given Y_T alone, the difference $A_l(T) - A_{l+1}(T)$ is stochastically dominated by Z_l. By the same argument as for (30), it follows that (31) is smaller than or equal to

$$\begin{aligned}\mathbb{E}\bigg(\sum_{l=2}^{Y_T-1} \sigma_S \frac{1}{\sqrt{l-1}} \frac{\beta}{(l-1)\lambda+\beta} \mathbb{1}_{\{J_T \leq l\}}\bigg) &= \mathbb{E}\bigg(\sum_{l=2}^{Y_T-1} \sigma_S \frac{1}{\sqrt{l-1}} \frac{\beta}{(l-1)\lambda+\beta} \mathbb{E}(\mathbb{1}_{\{J_T \leq l\}}|Y_T)\bigg) \\ &\leq \mathbb{E}\bigg(\sum_{l=2}^{Y_T-1} \sigma_S \frac{1}{\sqrt{l-1}} \frac{\beta}{(l-1)\lambda} \frac{l}{Y_T}\bigg) \\ &\leq \mathbb{E}\bigg(\sum_{l=2}^{Y_T-1} \sigma_S \frac{1}{\sqrt{l-1}} \frac{\beta}{(l-1)\lambda} \frac{2(l-1)}{Y_T}\bigg) \\ &\leq \frac{2\beta}{\lambda}\sigma_S \mathbb{E}\bigg(\frac{1}{Y_T}\sum_{l=2}^{Y_T-1} \frac{1}{\sqrt{l-1}}\bigg) \\ &\leq \frac{2\beta}{\lambda}\sigma_S \mathbb{E}\bigg(\frac{1}{Y_T}\sum_{l=2}^{Y_T-1} \frac{2}{\sqrt{l-2}+\sqrt{l-1}}\bigg) \\ &= \frac{2\beta}{\lambda}\sigma_S \mathbb{E}\bigg(\frac{1}{Y_T}\sum_{l=2}^{Y_T-1} 2(\sqrt{l-1}-\sqrt{l-2})\bigg) \\ &= \frac{4\beta}{\lambda}\sigma_S \mathbb{E}\bigg(\frac{\sqrt{Y_T-2}}{Y_T}\bigg) \\ &\leq \frac{4\beta}{\lambda}\sigma_S \mathbb{E}\bigg(\frac{1}{\sqrt{Y_T}}\bigg).\end{aligned}$$

Since we arranged $A_{J_T}(T) \leq A_{J_\infty}$, for the third line of the right-hand side of (29) we obtain

$$\begin{aligned}&\mathbb{E}\bigg|\mathbb{E}(S)(1-e^{-\beta A_{J_\infty}}) - \sum_{\substack{i=1\\ i \neq J_T}}^{Y_T} \sum_{l=i \vee J_T}^{Y_T-1} \frac{\mathbb{E}(S)}{l-1}(e^{-\beta A_{l+1}(T)} - e^{-\beta A_l(T)})\bigg| \\ &= \mathbb{E}\bigg|\mathbb{E}(S)(1-e^{-\beta A_{J_\infty}}) - \mathbb{E}(S)\sum_{l=J_T \vee 2}^{Y_T-1} \frac{1}{l-1} \sum_{\substack{i=1\\ i \neq J_T}}^{l} (e^{-\beta A_{l+1}(T)} - e^{-\beta A_l(T)})\bigg| \\ &= \mathbb{E}\bigg|\mathbb{E}(S)(1-e^{-\beta A_{J_\infty}}) - \mathbb{E}(S)\sum_{l=J_T \vee 2}^{Y_T-1} (e^{-\beta A_{l+1}(T)} - e^{-\beta A_l(T)})\bigg| \\ &= \mathbb{E}\bigg|\mathbb{E}(S)(1-e^{-\beta A_{J_\infty}}) - \mathbb{E}(S)(e^{-\beta A_{Y_T}(T)} - e^{-\beta A_{\max(J_T,2)}(T)})\bigg| \\ &= \mathbb{E}\bigg(\mathbb{E}(S)(1-e^{-\beta A_{Y_T}(T)}) + \mathbb{E}(S)(e^{-\beta A_{\max(J_T,2)}(T)} - e^{-\beta A_{J_\infty}})\bigg)\end{aligned}$$

$$= \mathbb{E}(S)\mathbb{E}\Big(1 - \mathbb{E}(e^{-\beta A_{Y_T}(T)}|Y_T)\Big) + \mathbb{E}(S)(\mathbb{E}(e^{-\beta A_{\max(J_T,2)}(T)}) - \mathbb{E}(e^{-\beta A_{J_\infty}})). \tag{32}$$

By Proposition 3.9

$$\mathbb{E}(e^{-\beta A_{\max(J_T,2)}(T)}) = \int_0^T e^{-\beta x}\frac{\lambda e^{-\lambda x}}{1-e^{-\lambda T}}dx = \frac{\lambda}{1-e^{-\lambda T}}\int_0^T e^{-(\beta+\lambda)x}dx = \frac{\lambda}{\beta+\lambda}\frac{1-e^{-(\beta+\lambda)T}}{1-e^{-\lambda T}}, \tag{33}$$

whence follows that (32) is smaller than or equal to

$$\mathbb{E}(S)\mathbb{E}\Big(\frac{\beta}{\lambda}\frac{1}{Y_t+\frac{\beta}{\lambda}}\Big) + \mathbb{E}(S)\frac{\lambda}{\beta+\lambda}\Big(\frac{1-e^{-(\beta+\lambda)T}}{1-e^{-\lambda T}} - 1\Big) \leq \frac{\beta}{\lambda}\mathbb{E}(S)\mathbb{E}\Big(\frac{1}{Y_T}\Big) + \mathbb{E}(S)\frac{\lambda}{\beta+\lambda}\frac{1}{e^{\lambda T}-1}.$$

Since we arranged $S_{J_T} = S_{J_\infty}$ and $A_{J_T}(T) \leq A_{J_\infty}$, for the fourth line of the right-hand side of (29) we have

$$\mathbb{E}\Big|S_{J_\infty}(1-e^{-\beta A_{J_\infty}}) - S_{J_T}(1-e^{-\beta A_{\max(J_T,2)}(T)})\Big| = \mathbb{E}(S)(\mathbb{E}(e^{-\beta A_{\max(J_T,2)}(T)}) - \mathbb{E}(e^{-\beta A_{J_\infty}})). \tag{34}$$

Note that the right-hand side of (34) is the same as the second summand of the right-hand side of (32), which is smaller than or equal to $\mathbb{E}(S)\frac{\lambda}{\beta+\mu}\frac{1}{e^{\lambda T}-1}$ (see above).

Altogether, we obtain the following upper bound for the right-hand side of (29):

$$\frac{2\alpha}{\lambda}\mathbb{E}(S)\mathbb{E}\Big(\frac{1}{Y_T}\Big) + \frac{4\alpha}{\lambda}\sigma_S\mathbb{E}\Big(\frac{1}{\sqrt{Y_T}}\Big) + \frac{2\alpha}{\beta}\mathbb{E}(S)\frac{\lambda}{\beta+\lambda}\frac{1}{e^{\lambda T}-1}.$$

□

5.2 The general case

5.2.1 The main theorem

Let $S_{J_\infty}(\omega) = S_{J_T}(\omega)$ as in the pure birth case. Furthermore, let $A_{J_\infty} = Z$, where Z is the random variable from Corollary 3.14. From the proof of this corollary, we see that

$$A_{J_\infty}(\omega) = \mathbb{1}_{\{J_T(\omega)<Y_T(\omega)\}}F_\infty^{-1}(F_*(A_{J_T}(T,\omega)) + \mathbb{1}_{\{J_T(\omega)=Y_T(\omega)\}}\tilde{Z}(\omega),$$

where F_∞ is the cumulative distribution function of the Exp(λ) distribution as before, $\tilde{Z} \sim F_\infty$ independent of everything and $F_*(t) = 1 - \frac{e^{-\lambda t} - e^{-(\lambda-\mu)T}e^{-\mu t}}{1-e^{-(\lambda-\mu)T}}$ (see also (16)). Note, given $Y_T > 0$, we have $A_{J_\infty} \sim$ Exp(λ).

Recall the definition of the parameter random variable Λ_T from (27) and let Λ_T^* be a random variable with $\mathcal{L}(\Lambda_T^*) = \mathcal{L}(\Lambda_T|Y_T > 0)$ as before. Moreover, set

$$\mathrm{M} := \frac{\alpha S_{J_\infty}}{\beta+\mu}(1-e^{-(\beta+\mu)A_{J_\infty}}) + \frac{\alpha\mathbb{E}(S)}{\beta+\mu}(1-e^{-(\beta+\mu)A_{J_\infty}}) \tag{35}$$

and let M^* be a random variable with $\mathcal{L}(\mathrm{M}^*) = \mathcal{L}(\mathrm{M}|Y_T > 0)$.

We already know that MixPo(Λ_T^*) is the degree distribution at time T in the general case. Theorem 5.4 below implies that MixPo(Λ_T^*) converges at a rate of just a bit over $e^{-\frac{1}{6}(\lambda-\mu)T}$ to the MixPo(M^*) distribution, which is the asymptotic degree distribution stated in Section 3.2 of [BL10], as $T \to \infty$. Note again that this theorem is much more powerful since it gives an exact bound for the total variation distance for every finite T.

5.4 Theorem
Let $\sigma_S < \infty$ be the standard deviation of S. Then for $T \geq \frac{2\log(4(\lambda(\lambda-\mu)^{-1}))}{\lambda-\mu}$, we have

$$
\begin{aligned}
&d_{TV}(\mathrm{MixPo}(\Lambda_T^*), \mathrm{MixPo}(\mathrm{M}^*)) \\
&\leq \alpha\left(\left(\frac{5\sqrt{6}}{2}\frac{\lambda}{\lambda-\mu} + \frac{2}{5} + \left(\beta + \frac{\mu}{2}\right)\left(\frac{229}{5(\lambda-\mu)} + \frac{2}{\lambda+\mu}\right)\right)\mathbb{E}(S) + \frac{27}{10}\sqrt{2}\sigma_S\right)(\lambda-\mu)T^2 e^{-\frac{1}{6}(\lambda-\mu)T} \\
&\quad + \alpha\beta\left(\left(\frac{1}{2} + \frac{\mu}{4\beta}\right)\frac{\mu}{\lambda} + \left(\frac{8}{\mu} + \frac{4}{\beta}\right)\frac{\lambda^3(\lambda+\mu)}{(\lambda-\mu)^3} + \left(6T + \frac{\sqrt{6}}{2} + \frac{3}{\beta}(\mu T + 3) + \frac{5\lambda}{\beta^2}\right)\lambda\right)\mathbb{E}(S)T^2 e^{-(\lambda-\mu)T} \\
&\quad + 3\sqrt{2}\alpha\lambda\sigma_S T^2 e^{-(\lambda-\mu)T}.
\end{aligned}
$$

Note that the right-hand side is of order

$$O(T^2)e^{-\frac{1}{6}(\lambda-\mu)T}$$

as $T \to \infty$.

The main idea of the proof of this theorem is the same as in the pure birth case: We make use of Theorem 2.1 to obtain $\mathbb{E}(|\Lambda_T - \mathrm{M}|\ |Y_T > 0)$ as an upper bound for $d_{TV}(\mathrm{MixPo}(\Lambda_T^*), \mathrm{MixPo}(\mathrm{M}^*))$ and establish a further bound for this expected value. In order to do so, we use that the expected value of the first summand of the right-hand side of (27) converges quickly to zero since the time $T - T_{B_T+D_T}$ since the last event before T converges quickly to 0, and compare the remaining summand of the right-hand side of (27) with the right-hand side of (35). For this comparison, we make vital use of the fact that the average of the social indices of the nodes living at time T is close to $\mathbb{E}(S)$ by the Law of Large Numbers again and that, given T_l for some large $l \in \mathbb{N}$, the percentage of the nodes living at time T_l that survive up to time T is approximately $e^{-\mu(T-T_l)}$ by the Law of Large Numbers (see Lemma 5.17).

A further important ingredient is that the reciprocal of the node process $(Y_t)_{t\geq 0}$ conditioned on survival is a supermartingale (see Corollary 5.8), which makes it easy to deal with the expected value of its maximum. Finally, we also use that the age $A_{J_T}(T)$ of the randomly picked individual converges quickly to A_{J_∞} by Corollary 3.14.

The fact that nodes may die complicates the procedure considerably since the population size after a fixed number of events is random in this case and additional dependencies have to be treated (e.g. the inter-event times depend on the random population size at the previous event time).

In order to cope with additional dependencies, we introduce a deterministic number $\kappa(T)$ that depends on T such that the probability for more than $\kappa(T)$ events up to time T decreases exponentially in T (see Definition 5.9 and Lemma 5.10(ii) below). Essentially, we substitute the number of events $B_T + D_T$ up to time T by $\kappa(T)$ in the proof of Lemma 5.19 below since it is difficult to treat the dependencies between the number of events up to time T and the event times.

In order to cope with additional dependencies on the random index J_T, we note that the probability that the node that is randomly picked at time T was born up to time $\frac{T}{2}$ decreases exponentially in T (see Lemma 5.10(i)). Thus we essentially only have to consider the time interval $[\frac{T}{2}, \infty)$ instead of the interval $[T_{r(J_T)}, \infty)$, where $T_{r(J_T)}$ is the birth time of the randomly picked individual. The choice of $\frac{T}{2}$ as the left endpoint of the interval makes sure that we always have a large number of individuals in the time interval with high probability.

5.2.2 Some lemmas of general interest

Before we prove Theorem 5.4 in detail, we formulate several important statements that are used in the proof and could also be useful in other settings. The first one gives us an expression for the extinction probability given that the process survives up to time T. In order to simplify the notation, we define $Y_\infty := \lim\limits_{t \to \infty} Y_t \in \{0, \infty\}$.

5.5 Lemma
For the conditioned extinction probability given $Y_T > 0$, we have

$$\mathbb{P}(Y_\infty = 0 | Y_T > 0) = \frac{\mu}{\lambda} e^{-(\lambda-\mu)T}.$$

Proof: By conditioning on the population size, we obtain

$$\mathbb{P}(Y_\infty = 0 | Y_T > 0) = \mathbb{E}(\mathbb{P}(Y_\infty = 0 | Y_T) | Y_T > 0) = \mathbb{E}\left(\left(\frac{\mu}{\lambda} \right)^{Y_T} \middle| \, Y_T > 0 \right) \tag{36}$$

since $(\mu/\lambda)^m$ is the extinction probability of a linear birth and death process with birth rate λ, death rate μ and initial value m (see Remark 3.1).

On the one hand, we have

$$\begin{aligned}\mathbb{E}\left(\left(\frac{\mu}{\lambda} \right)^{Y_T} \right) &= \mathbb{E}\left(\left(\frac{\mu}{\lambda} \right)^{Y_T} \middle| \, Y_T > 0 \right) \mathbb{P}(Y_T > 0) + \mathbb{E}\left(\left(\frac{\mu}{\lambda} \right)^{Y_T} \middle| \, Y_T = 0 \right) \mathbb{P}(Y_T = 0) \\ &= \mathbb{E}\left(\left(\frac{\mu}{\lambda} \right)^{Y_T} \middle| \, Y_T > 0 \right) (1 - p_0(T)) + p_0(T).\end{aligned}$$

On the other hand, we can make use of the known formula for the probability generating function of Y_T (see III.5 in [AN72]) to obtain

$$\mathbb{E}\left(\left(\frac{\mu}{\lambda} \right)^{Y_T} \right) = \frac{\mu}{\lambda}.$$

This yields

$$\mathbb{E}\left(\left(\frac{\mu}{\lambda} \right)^{Y_T} \middle| \, Y_T > 0 \right) = \left(\frac{\mu}{\lambda} - p_0(T) \right) (1 - p_0(T))^{-1} = \frac{\mu(\lambda - \mu)}{\lambda(\lambda e^{(\lambda-\mu)T} - \mu)} \frac{\lambda e^{(\lambda-\mu)T} - \mu}{(\lambda - \mu) e^{(\lambda-\mu)T}} \leq \frac{\mu}{\lambda} e^{-(\lambda-\mu)}.$$

□

For the probability of $Y_T = 1$ conditioned on $Y_T > 0$, we have the following lemma.

5.6 Lemma
We have for $T \geq \frac{1}{\lambda-\mu} \log(2)$

$$\mathbb{P}(Y_T = 1 | Y_T > 0) \leq \frac{2(\lambda - \mu)}{\lambda e^{(\lambda-\mu)T}}.$$

Proof: Using the probability mass function p from Subsection 3.1, for $T \geq \frac{1}{\lambda-\mu}\log(2)$, we obtain

$$\mathbb{P}(Y_T = 1 | Y_T > 0) = \frac{p_1(T)}{1-p_0(T)} = 1 - \lambda\tilde{p}(T) = \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)T} - \mu} \leq \frac{2(\lambda-\mu)}{\lambda e^{(\lambda-\mu)T}}.$$

□

We next consider sub- and supermartingal properties of conditioned processes.

5.7 Lemma

(i) $(Y_t)_{t\geq 0}$ conditioned on $Y_\infty = 0$ is a supermartingale.

(ii) $(Y_t)_{t\geq 0}$ conditioned on $Y_\infty > 0$ is a submartingale.

Proof:

(i) Consider a subcritical linear birth and death process $(\tilde{Y}_t)_{t\geq 0}$ with *birth rate* μ, *death rate* λ and initial value one. Then $(\tilde{Y}_t)_{t\geq 0}$ has the same law as $(Y_t)_{t\geq 0}$ conditioned on $Y_\infty = 0$ (see e.g. page 78 in [Lam08]), which can be proved by using the embedded generation process (see e.g. Section IV.3 of [AN72]) and the corresponding result for discrete-time branching processes (see e.g. Theorem 3 in Section I.12 of [AN72]). It is well-known that a subcritical linear birth and death process is a submartingale, which yields the result.

(ii) Consider a process $(\hat{Y}_t)_{t\geq 0}$ that has the law of $(Y_t)_{t\geq 0}$ conditioned on $Y_\infty > 0$. Note that $(\hat{Y}_t)_{t\geq 0}$ inherits the Markov property from $(Y_t)_{t\geq 0}$. Furthermore, it is well-known that $(Y_t)_{t\geq 0}$ is a submartingale. Thus we obtain for $t > s \geq 0$ and $y_s \in \mathbb{N}$

$$\begin{aligned}
\mathbb{E}(\hat{Y}_t | \hat{Y}_s = y_s) &= \mathbb{E}(Y_t | Y_s = y_s, Y_\infty > 0) \\
&= \frac{\mathbb{E}(\mathbb{1}_{\{Y_\infty > 0\}} Y_t | Y_s = y_s)}{\mathbb{P}(Y_\infty > 0 | Y_s = y_s)} \\
&= \frac{\mathbb{E}(Y_t | Y_s = y_s) - \mathbb{E}(\mathbb{1}_{\{Y_\infty = 0\}} Y_t | Y_s = y_s)}{\mathbb{P}(Y_\infty > 0 | Y_s = y_s)} \\
&\geq \frac{y_s - \mathbb{P}(Y_\infty = 0 | Y_s = y_s)\mathbb{E}(\tilde{Y}_t | \tilde{Y}_s = y_s)}{\mathbb{P}(Y_\infty > 0 | Y_s = y_s)} \\
&\geq \frac{y_s - \mathbb{P}(Y_\infty = 0 | Y_s = y_s) y_s}{\mathbb{P}(Y_\infty > 0 | Y_s = y_s)} \\
&= y_s,
\end{aligned}$$

where $(\tilde{Y}_t)_{t\geq 0}$ is the supermartingale from (i), which also inherits the Markov property from $(Y_t)_{t\geq 0}$. Thus $(Y_t)_{t\geq 0}$ conditioned on ultimate survival is a submartingale. □

Lemma 5.7 yields a useful result about $(Y_t^{-1})_{t\geq 0}$ conditioned on ultimate survival:

5.8 Corollary
$(Y_t^{-1})_{t\geq 0}$ conditioned on $Y_\infty > 0$ is a supermartingale.

Proof: In general, for a submartingale $(Z_t)_{t\geq 0}$ with respect to a filtration $(\mathcal{F}_t)_{t\geq 0}$ with $Z_t \geq 1$ for all $t \geq 0$, we have for $t > s \geq 0$

$$\mathbb{E}\biggl(\frac{1}{Z_t} - \frac{1}{Z_s} \biggm| \mathcal{F}_s\biggr) = \mathbb{E}\biggl(\frac{Z_s - Z_t}{Z_s Z_t} \biggm| \mathcal{F}_s\biggr) \leq \mathbb{E}(Z_s - Z_t|\mathcal{F}_s) \leq 0.$$

Thus $(Z_t^{-1})_{t\geq 0}$ is a supermartingale. Consequently, Lemma 5.7 implies that $(Y_t^{-1})_{t\geq 0}$ conditioned on ultimate survival is a supermartingale. ☐

Now we introduce the deterministic number $\kappa(T)$ as mentioned before, which helps us to cope with dependencies. Furthermore, we define the random number $\mathcal{K}(T)$ as index of the last event time before $\frac{T}{2}$ such that $r(J_T) > \mathcal{K}(T)$ is equivalent to $T_{r(J_T)} \geq \frac{T}{2}$ and $\mathcal{M}_T$ as the sum of the number of births and the number of deaths to simplify notation.

5.9 Definition
Let $\kappa(T) := \lfloor e^{\frac{3}{2}(\lambda+\mu)T} \rfloor$ and $\mathcal{K}(T) := \max\{k : T_k < \frac{T}{2}\}$, such that $Y_{T_{\mathcal{K}(T)}} = Y_{\frac{T}{2}}$ almost surely. Moreover, let $\mathcal{M}_T := B_T + D_T$ be the number of events up to time T.

The following lemma implies the properties of $\kappa(T)$ and $\mathcal{K}(T)$ that were stated before and that we need to prove the convergence rate for the degree distribution in the general case.

5.10 Lemma

(i) For the probability that fewer than $\mathcal{K}(T)$ events have occurred up to the birth time of the randomly picked node J_T was born, we have

$$\begin{aligned}\mathbb{P}(r(J_T) \leq \mathcal{K}(T)|Y_T > 0) &= \mathbb{P}\biggl(T_{r(J_T)} < \frac{T}{2} \biggm| Y_T > 0\biggr)\\ &\leq e^{-\frac{1}{2}\lambda T} + \frac{\lambda - \mu}{\lambda e^{(\lambda-\mu)T} - \lambda}\biggl(\log\biggl(\frac{\lambda}{\lambda - \mu}\biggr) + (\lambda - \mu)T\biggr).\end{aligned}$$

(ii) For $T \geq \frac{2(\log(4\lambda) - \log(\lambda-\mu))}{\lambda+\mu}$, we have

$$\mathbb{P}(\kappa(T) \leq \mathcal{M}_T|Y_T > 0) \leq \frac{60\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}e^{-(\lambda+\mu)T}$$

Proof:

(i) By Remark 3.17, we have

$$\begin{aligned}\mathbb{P}(r(J_T) \leq \mathcal{K}(T)|Y_T > 0) = \mathbb{P}(T_{r(J_T)} \leq T_{\mathcal{K}(T)}|Y_T > 0) &= \mathbb{P}\biggl(T - T_{r(J_T)} \geq \frac{T}{2} \biggm| Y_T > 0\biggr)\\ \leq \mathbb{E}\biggl(\frac{Y_T - 1}{Y_T}e^{-\frac{1}{2}\lambda T} + \frac{1}{Y_T}\biggm|Y_T > 0\biggr) &\leq e^{-\frac{1}{2}\lambda T} + \mathbb{E}\biggl(\frac{1}{Y_T}\biggm|Y_T > 0\biggr). \quad (37)\end{aligned}$$

By Proposition 3.3, the right-hand side of (37) is smaller than or equal to

$$e^{-\frac{1}{2}\lambda T} + \frac{\lambda - \mu}{\lambda e^{(\lambda-\mu)T} - \lambda}\biggl(\log\biggl(\frac{\lambda}{\lambda - \mu}\biggr) + (\lambda - \mu)T\biggr).$$

(ii) Using Proposition 3.6, we compute

$$\begin{aligned}\kappa(T) - 2\mathbb{E}(B_T) &\geq e^{\frac{3}{2}(\lambda+\mu)T} - 1 - \frac{2\lambda}{\lambda-\mu}e^{(\lambda-\mu)T} + \frac{2\mu}{\lambda-\mu}\\ &= e^{(\lambda-\mu)T}\left(e^{2\mu T}e^{\frac{1}{2}(\lambda+\mu)T} - \frac{2\lambda}{\lambda-\mu}\right) + \frac{3\mu-\lambda}{\lambda-\mu}. \qquad (38)\end{aligned}$$

Since $T \geq \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu}$, we have

$$\frac{1}{2}e^{2\mu T}e^{\frac{1}{2}(\lambda+\mu)T} \geq \frac{2\lambda}{\lambda-\mu}. \qquad (39)$$

Thus for $T \geq \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu}$, the right-hand side of (38) is larger than or equal to

$$e^{(\lambda-\mu)T}\frac{1}{2}e^{2\mu T}e^{\frac{1}{2}(\lambda+\mu)T} + \frac{3\mu-\lambda}{\lambda-\mu}. \qquad (40)$$

Since for $T \geq \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu}$, we have

$$e^{(\lambda-\mu)T} \geq 1 + (\lambda-\mu)T \geq 1 + 2\log\left(\frac{4\lambda}{\lambda+\mu}\right) \geq 1 + 2\log(4) \geq 2, \qquad (41)$$

inequality (39) implies that for $T \geq \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda-\mu}$, the expression (40) is bounded from below by

$$e^{(\lambda-\mu)T}\frac{3}{8}e^{2\mu T}e^{\frac{1}{2}(\lambda+\mu)T} + \frac{3\mu}{\lambda-\mu} \geq \frac{3}{8}e^{\frac{3}{2}(\lambda+\mu)T}.$$

Thus for $T \geq \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu}$, we have

$$\kappa(T) - 2\mathbb{E}(B_T) \geq \frac{3}{8}e^{\frac{3}{2}(\lambda+\mu)T} > 0. \qquad (42)$$

Consequently, we can apply Chebyshev's inequality and obtain

$$\begin{aligned}\mathbb{P}(\kappa(T) \leq \mathcal{M}_T | Y_T > 0) &\leq \frac{\mathbb{P}(\kappa(T) \leq \mathcal{M}_T)}{\mathbb{P}(Y_T > 0)} \leq \frac{\mathbb{P}(\kappa(T) \leq 2B_T)}{\mathbb{P}(Y_T > 0)}\\ &= \frac{1}{\mathbb{P}(Y_T > 0)}\mathbb{P}(\kappa(T) - 2\mathbb{E}(B_T) \leq 2B_T - 2\mathbb{E}(B_T))\\ &\leq \frac{1}{\mathbb{P}(Y_T > 0)}\frac{4\mathrm{Var}(B_T)}{(\kappa(T) - 2\mathbb{E}(B_T))^2}, \qquad (43)\end{aligned}$$

where $T \geq \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu}$. By $\mathbb{P}(Y_T > 0) \geq \mathbb{P}(Y_\infty > 0) = \frac{\lambda-\mu}{\lambda}$, Proposition 3.6 and (42), for $T \geq \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu}$, the right-hand side of (43) is smaller than or equal to

$$\frac{256}{9}\frac{\lambda}{\lambda-\mu}\frac{\mathrm{Var}(B_T)}{e^{3(\lambda+\mu)T}} \leq \frac{30\lambda}{\lambda-\mu}\left(\frac{\lambda^2(\lambda+\mu)}{(\lambda-\mu)^3}e^{-(\lambda+\mu)T} + \frac{2\lambda^2\mu}{(\lambda-\mu)^3}e^{-2(\lambda+\mu)T}\right). \qquad (44)$$

Since $e^{-(\lambda+\mu)T} \leq \frac{1}{2}$ by (41), the right-hand side of (44) is smaller than or equal to

$$\frac{60\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}e^{-(\lambda+\mu)T}.$$

□

5.2.3 Further lemmas

Now we give further statements that are used in order to prove Theorem 5.4.

The following lemma states that the time $T - T_{\mathcal{M}_T}$ since the last event becomes small quickly.

5.11 Lemma

For $T \geq \frac{1}{\lambda-\mu}\log(2)$ and $c > 0$, we have

$$\mathbb{E}(1 - e^{-c(T-T_{\mathcal{M}_T})}|Y_T > 0) \leq \frac{2(\lambda-\mu)}{\lambda e^{(\lambda-\mu)T}} + \frac{c}{\lambda}\mathbb{E}\bigg(\frac{1}{Y_T - 1}\mathbb{1}_{\{Y_T>1\}}\ \bigg|\ Y_T > 0\bigg). \tag{45}$$

Proof: Let X have the cumulative distribution function

$$G(t) = \mathbb{1}_{\{Y_T>1\}}(1 - e^{-(Y_T-1)(\lambda-\mu)t}) + \mathbb{1}_{\{Y_T\leq1\}}\mathbb{1}_{\{t\geq T\}}.$$

Conditionally on Y_T, we then have $T - T_{\mathcal{M}_T} \leq_{st} X$ by Theorem 3.19, which implies

$$\begin{aligned}\mathbb{E}(1-&e^{-c(T-T_{\mathcal{M}_T})}|Y_T > 0) \leq \mathbb{E}(1 - e^{-cX}|Y_T > 0)\\ &= \mathbb{E}\bigg((1 - e^{-cT})\mathbb{P}(X = T|Y_T) + \mathbb{E}(1 - e^{-cX}|X < T, Y_T)\mathbb{P}(X < T|Y_T)\ \bigg|\ Y_T > 0\bigg). \end{aligned}\tag{46}$$

Since $\mathbb{P}(X = T|Y_T = 1) = 1$ and $\mathcal{L}(X|Y_T) = \mathrm{Exp}((Y_T - 1)\lambda)$ on $\{Y_T \geq 2\}$, we obtain that the right-hand side of (46) is smaller than or equal to

$$\begin{aligned}\mathbb{E}\bigg(\mathbb{1}_{\{Y_T=1\}} + \frac{c}{c + (Y_T - 1)\lambda}\mathbb{1}_{\{Y_T>1\}}\ \bigg|\ Y_T > 0\bigg) &\leq \mathbb{P}(Y_T = 1|Y_T > 0)\\ &\quad + \frac{c}{\lambda}\mathbb{E}\bigg(\frac{1}{Y_T - 1}\mathbb{1}_{\{Y_T>1\}}\ \bigg|\ Y_T > 0\bigg).\end{aligned}$$

For the conditional probability of $Y_T = 1$, by Lemma 5.6, we have for $T \geq \frac{1}{\lambda-\mu}\log(2)$

$$\mathbb{P}(Y_T = 1|Y_T > 0) \leq \frac{2(\lambda-\mu)}{\lambda e^{(\lambda-\mu)T}}.$$

□

The following lemma gives us an upper bound for the conditional expectation on the right-hand side of (45) and is proved similarly to Proposition 3.3.

5.12 Lemma

We have

$$\mathbb{E}\bigg(\frac{1}{Y_T - 1}\mathbb{1}_{\{Y_T>1\}}\ \bigg|\ Y_T > 0\bigg) \leq \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)T} - \mu}\bigg(\log\bigg(\frac{\lambda}{\lambda-\mu}\bigg) + (\lambda-\mu)T\bigg).$$

Proof: We have

$$\mathbb{E}\bigg(\frac{1}{Y_T - 1}\mathbb{1}_{\{Y_T>1\}}\ \bigg|\ Y_T > 0\bigg) = \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)T} - \mu}\sum_{n=1}^{\infty}\frac{1}{n}\bigg(\frac{\lambda e^{(\lambda-\mu)T} - \lambda}{\lambda e^{(\lambda-\mu)T} - \mu}\bigg)^n \tag{47}$$

and in the proof of Proposition 3.3, we computed an upper bound for the series in (47) that yields that the right-hand side of (47) is smaller than or equal to

$$\frac{\lambda-\mu}{\lambda e^{-(\lambda-\mu)T} - \mu}\bigg(\log\bigg(\frac{\lambda}{\lambda-\mu}\bigg) + (\lambda-\mu)T\bigg).$$

□

In order to prove our next lemma, we use the supermartingale from Corollary 5.8.

5.13 Lemma
For all $\delta, \gamma > 0$, we have

$$\mathbb{E}\left(\frac{1}{\min\limits_{\frac{T}{2}\leq t} Y_t^\delta} \,\middle|\, Y_\infty > 0\right) \leq e^{-\gamma(\lambda-\mu)T} + \mathbb{E}\left(\frac{1}{Y_{\frac{T}{2}}} \,\middle|\, Y_\infty > 0\right) e^{\frac{\gamma}{\delta}(\lambda-\mu)T}.$$

Proof: We obtain

$$\begin{aligned}\mathbb{E}\left(\frac{1}{\min\limits_{\frac{T}{2}\leq t} Y_t^\delta} \,\middle|\, Y_\infty > 0\right) &\leq \mathbb{E}\left(\mathbb{E}\left(\mathbb{1}_{\{\max\limits_{\frac{T}{2}\leq t} Y_t^{-\delta} \leq e^{-\gamma(\lambda-\mu)T}\}} \max_{\frac{T}{2}\leq t} \frac{1}{Y_t^\delta} \,\middle|\, Y_{\frac{T}{2}}\right) \,\middle|\, Y_\infty > 0\right) \\ &\quad + \mathbb{E}\left(\mathbb{E}\left(\mathbb{1}_{\{\max\limits_{\frac{T}{2}\leq t} Y_t^{-\delta} > e^{-\gamma(\lambda-\mu)T}\}} \max_{\frac{T}{2}\leq t} \frac{1}{Y_t^\delta} \,\middle|\, Y_{\frac{T}{2}}\right) \,\middle|\, Y_\infty > 0\right) \\ &\leq e^{-\gamma(\lambda-\mu)T} + \mathbb{E}\left(\mathbb{P}\left(\max_{\frac{T}{2}\leq t} \frac{1}{Y_t} > e^{-\frac{\gamma}{\delta}(\lambda-\mu)T} \,\middle|\, Y_{\frac{T}{2}}\right) \,\middle|\, Y_\infty > 0\right) \\ &\leq e^{-\gamma(\lambda-\mu)T} + \mathbb{E}\left(\frac{1}{Y_{\frac{T}{2}}} \,\middle|\, Y_\infty > 0\right) e^{\frac{\gamma}{\delta}(\lambda-\mu)T},\end{aligned}$$

where the last line follows from Corollary 5.8 and the the inequality (2) in Theorem 6.14 on page 99 in [Yeh95]. ☐

We use the following lemma to bound the conditional expectation in the second summand of the right-hand side of Lemma 5.13 from above.

5.14 Lemma
We have for $T \geq \frac{2}{\lambda-\mu}\log(2)$

$$\mathbb{E}\left(\frac{1}{Y_{\frac{T}{2}}} \,\middle|\, Y_\infty > 0\right) \leq 2\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + \frac{1}{2}(\lambda-\mu)T\right) e^{-\frac{1}{2}(\lambda-\mu)T}.$$

Proof: For $T \geq \frac{2}{\lambda-\mu}\log(2)$, we obtain

$$\begin{aligned}\mathbb{E}\left(\frac{1}{Y_{\frac{T}{2}}} \,\middle|\, Y_\infty > 0\right) &= \frac{\mathbb{E}\left(Y_{\frac{T}{2}}^{-1}\mathbb{1}_{\{Y_\infty>0\}}\right)}{\mathbb{P}(Y_\infty > 0)} \leq \frac{\lambda}{\lambda-\mu}\mathbb{E}\left(Y_{\frac{T}{2}}^{-1}\mathbb{1}_{\{Y_{\frac{T}{2}}>0\}}\right) \leq \frac{\lambda}{\lambda-\mu}\mathbb{E}\left(\frac{1}{Y_{\frac{T}{2}}} \,\middle|\, Y_{\frac{T}{2}} > 0\right) \\ &\leq \frac{\lambda}{\lambda e^{\frac{1}{2}(\lambda-\mu)T} - \lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + \frac{1}{2}(\lambda-\mu)T\right) \\ &\leq \frac{1}{\frac{1}{2}e^{\frac{1}{2}(\lambda-\mu)T} + \frac{1}{2}e^{\frac{1}{2}(\lambda-\mu)T} - 1}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + \frac{1}{2}(\lambda-\mu)T\right) \\ &\leq 2\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + \frac{1}{2}(\lambda-\mu)T\right) e^{-\frac{1}{2}(\lambda-\mu)T},\end{aligned}$$

where the second line follows from Proposition 3.3 and the last inequality holds for $T \geq \frac{2}{\lambda-\mu}\log(2)$. ☐

Before we continue to state and prove further useful statements, we introduce another random variable.

5.15 Definition
Let $R_{T_l,T}$ be the number of nodes that are alive at time T_l and survive up to time T.

The following lemma gives us the first two conditional moments of $R_{T_l,T}$.

5.16 Lemma
We have

$$\begin{aligned}\mathbb{E}(R_{T_l,T}|Y_{T_l},T_l,T_{l+1}) &= (Y_{T_l}-1)e^{-\mu(T-T_{l+1})}\\ \mathbb{E}(R^2_{T_l,T}|Y_{T_l},T_l,T_{l+1}) &= Y_{T_l}e^{-\mu(T-T_{l+1})} - Y_{T_l}e^{-2\mu(T-T_{l+1})} + Y^2_{T_l}e^{-2\mu(T-T_{l+1})}\\ &\quad - \mathbb{P}(Y_{T_{l+1}} = Y_{T_l}-1|Y_{T_l},T_l,T_{l+1})\Big(2(Y_{T_l}-1)e^{-2\mu(T-T_{l+1})} + e^{-\mu(T-T_{l+1})}\Big).\end{aligned}$$

Proof: Firstly, we determine the conditional expectation of $R_{T_l,T}$:

$$\begin{aligned}\mathbb{E}(R_{T_l,T}|Y_{T_l},T_l,T_{l+1}) &= p^+\mathbb{E}(R_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}+1)\\ &\quad + p^-\mathbb{E}(R_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}-1),\end{aligned}\tag{48}$$

where $p^+ := \mathbb{P}(Y_{T_{l+1}} = Y_{T_l}+1|Y_{T_l},T_l,T_{l+1})$ and $p^- := \mathbb{P}(Y_{T_{l+1}} = Y_{T_l}-1|Y_{T_l},T_l,T_{l+1})$.[2]

With

$$\begin{aligned}\mathbb{E}(R_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}+1) &= \mathbb{E}(R_{T_{l+1},T} - \mathbb{1}_{\{T^-_{r^{-1}(l+1)}>T\}}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}+1)\\ &= (Y_{T_l}+1)e^{-\mu(T-T_{l+1})} - e^{-\mu(T-T_{l+1})} = Y_{T_l}e^{-\mu(T-T_{l+1})}\end{aligned}\tag{49}$$

and

$$\begin{aligned}\mathbb{E}(R_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}-1) &= \mathbb{E}(R_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}-1).\\ &= (Y_{T_l}-1)e^{-\mu(T-T_{l+1})}.\end{aligned}$$

Equation (48) implies

$$\mathbb{E}(R_{T_l,T}|Y_{T_l},T_l,T_{l+1}) = Y_{T_l}e^{-\mu(T-T_{l+1})} - e^{-\mu(T-T_{l+1})}p^-.$$

Secondly, we compute the conditional second moment of $R_{T_l,T}$:

$$\begin{aligned}\mathbb{E}(R^2_{T_l,T}|Y_{T_l},T_l,T_{l+1}) &= p^+\mathbb{E}(R^2_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}+1)\\ &\quad + p^-\mathbb{E}(R^2_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}-1).\end{aligned}\tag{50}$$

We treat the summands separately again. For the case where a birth occurs at time T_{l+1}, we have

$$\begin{aligned}&\mathbb{E}(R^2_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}+1)\\ &= \mathbb{E}((R_{T_{l+1},T} - \mathbb{1}_{\{T^-_{r^{-1}(l+1)}>T\}})^2|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}+1)\\ &= \mathbb{E}\Bigg(R^2_{T_{l+1},T} - 2R_{T_{l+1},T}\mathbb{1}_{\{T^-_{r^{-1}(l+1)}>T\}} + \mathbb{1}_{\{T^-_{r^{-1}(l+1)}>T\}}\ \Bigg|\ Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}} = Y_{T_l}+1,Y_{T_l}>0\Bigg)\end{aligned}\tag{51}$$

---

[2] We have $p^+ = \frac{\lambda}{\lambda+\mu}$ and $p^- = \frac{\mu}{\lambda+\mu}$. However, we do not need these explicit expressions for our purposes.

and further

$$\begin{aligned}\mathbb{E}(R^2_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1) &= \mathrm{Var}(R_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)\\ &\quad+(\mathbb{E}(R_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1))^2. \qquad (52)\end{aligned}$$

Given $Y_{T_{l+1}}$, let $\mathcal{L}_{l+1}$ be the set of the $Y_{T_{l+1}}$ nodes living at time T_{l+1}. Then by independence of various death times, we obtain for the conditional variance

$$\begin{aligned}&\mathrm{Var}(R_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)\\ &=\mathrm{Var}\biggl(\sum_{j\in\mathcal{L}_{l+1}}\mathbb{1}_{\{T^-_j>T\}}\biggm| Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1,Y_{T_l}>0\biggr)\\ &=(Y_{T_l}+1)(e^{-\mu(T-T_{l+1})}-e^{-2\mu(T-T_{l+1})}).\end{aligned}$$

For the second summand of (52), we obtain:

$$\mathbb{E}(R_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)=(Y_{T_l}+1)e^{-\mu(T-T_{l+1})}.$$

Thus (52) is equal to

$$(Y_{T_l}+1)\biggl(e^{-\mu(T-T_{l+1})}+Y_{T_l}e^{-2\mu(T-T_{l+1})}\biggr).$$

For the remaining parts of (51), we have

$$\begin{aligned}&\mathbb{E}(R_{T_{l+1},T}\mathbb{1}_{\{T^-_{r^{-1}(l+1)}>T\}}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)\\ &=\mathbb{P}(T^-_{r^{-1}(l+1)}>T|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)\\ &\quad\cdot\mathbb{E}(R_{T_{l+1},T}|T^-_{r^{-1}(l+1)}>T,Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)\\ &=e^{-\mu(T-T_{l+1})}(1+Y_{T_l}e^{-\mu(T-T_{l+1})})\end{aligned}$$

and

$$\mathbb{E}(\mathbb{1}_{\{T^-_{r^{-1}(l+1)}>T\}}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)=e^{-\mu(T-T_{l+1})}.$$

Thus (51) implies

$$\begin{aligned}&\mathbb{E}(R^2_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}+1)\\ &=(Y_{T_l}+1)\biggl(e^{-\mu(T-T_{l+1}}+Y_{T_l}e^{-2\mu(T-T_{l+1})}\biggr)-2\biggl(e^{-\mu(T-T_{l+1}}+Y_{T_l}e^{-2\mu(T-T_{l+1})}\biggr)+e^{-\mu(T-T_{l+1})}\\ &=Y_{T_l}e^{-\mu(T-T_{l+1})}-Y_{T_l}e^{-2\mu(T-T_{l+1})}+Y^2_{T_l}e^{-2\mu(T-T_{l+1})}.\end{aligned}$$

For the case where a death occurs at time T_{l+1}, we have

$$\begin{aligned}&\mathbb{E}(R^2_{T_l,T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}-1)\\ &=\mathrm{Var}(R_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}-1)+(\mathbb{E}(R_{T_{l+1},T}|Y_{T_l},T_l,T_{l+1},Y_{T_{l+1}}=Y_{T_l}-1))^2\\ &=(Y_{T_l}-1)(e^{-\mu(T-T_{l+1})}-e^{-2\mu(T-T_{l+1})})+(Y_{T_l}-1)^2e^{-2\mu(T-T_{l+1})}\end{aligned}$$

$$= (Y_{T_l} - 1)e^{-\mu(T-T_{l+1})} + (Y_{T_l}^2 - 3Y_{T_l} + 2)e^{-2\mu(T-T_{l+1})}.$$

Thus from (50) follows

$$\begin{aligned}\mathbb{E}(R_{T_l,T}^2|Y_{T_l}, T_l, T_{l+1}, Y_{T_l} > 0) = &Y_{T_l}e^{-\mu(T-T_{l+1})} - Y_{T_l}e^{-2\mu(T-T_{l+1})} + Y_{T_l}^2e^{-2\mu(T-T_{l+1})} \\ &+ p^-\left(2(1 - Y_{T_l})e^{-2\mu(T-T_{l+1})} - e^{-\mu(T-T_{l+1})}\right).\end{aligned}$$

□

Knowing the conditional moments of $R_{T_l,T}$, we can find an upper bound for a more complex conditional expectation involving $R_{T_l,T}$ that appears in the proof of the main theorem below:

5.17 Lemma

For $l \in \mathbb{N}$, we have

$$\mathbb{E}\left(\left|\frac{R_{T_l,T} - 1}{Y_{T_l} - 1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})}\right| \,\middle|\, Y_{T_l}, T_l, T_{l+1}\right) \leq \left(\frac{6}{Y_{T_l}}\right)^{\frac{1}{2}}.$$

Proof: By Jensen's inequality, we obtain

$$\begin{aligned}&\mathbb{E}\left(\left|\frac{R_{T_l,T} - 1}{Y_{T_l} - 1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})}\right| \,\middle|\, Y_{T_l}, T_l, T_{l+1}\right) \\ &\leq \left(\mathbb{E}\left(\left(\frac{R_{T_l,T} - 1}{Y_{T_l} - 1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})}\right)^2 \,\middle|\, Y_{T_l}, T_l, T_{l+1}\right)\right)^{\frac{1}{2}} \\ &= \left(\mathbb{1}_{\{Y_{T_l}>1\}}\frac{\mathbb{E}((R_{T_l,T} - 1)^2|Y_{T_l}, T_l, T_{l+1})}{(Y_{T_l} - 1)^2} - \frac{2}{Y_{T_l} - 1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\mu(T-T_{l+1})}\mathbb{E}(R_{T_l,T} - 1|Y_{T_l}, T_l, T_{l+1}) \right. \\ &\quad \left. + e^{-2\mu(T-T_{l+1})}\right)^{\frac{1}{2}}. \end{aligned} \tag{53}$$

Lemma 5.16 implies

$$\begin{aligned}\frac{2}{Y_{T_l} - 1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\mu(T-T_{l+1})}\mathbb{E}(R_{T_l,T} - 1|Y_{T_l}, T_l, T_{l+1}) =& 2 \cdot \mathbb{1}_{\{Y_{T_l}>1\}}e^{-2\mu(T-T_{l+1})} \\ &- \frac{2}{Y_{T_l} - 1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\mu(T-T_{l+1})}.\end{aligned} \tag{54}$$

and

$$\begin{aligned}&\frac{1}{(Y_{T_l} - 1)^2}\mathbb{1}_{\{Y_{T_l}>1\}}\mathbb{E}((R_{T_l,T} - 1)^2|Y_{T_l}, T_l, T_{l+1}) \\ &= \frac{1}{(Y_{T_l} - 1)^2}\mathbb{1}_{\{Y_{T_l}>1\}}\left(\mathbb{E}(R_{T_l,T}^2|Y_{T_l}, T_l, T_{l+1}) - 2\mathbb{E}(R_{T_l,T}|Y_{T_l}, T_l, T_{l+1}) + 1\right) \\ &= \frac{1}{(Y_{T_l} - 1)^2}\mathbb{1}_{\{Y_{T_l}>1\}}\left(Y_{T_l}e^{-\mu(T-T_{l+1})} - Y_{T_l}e^{-2\mu(T-T_{l+1})} + Y_{T_l}^2e^{-2\mu(T-T_{l+1})}\right. \\ &\qquad \left. - \mathbb{P}(Y_{T_{l+1}} = Y_{T_l} - 1|Y_{T_l}, T_l, T_{l+1})\left(2(Y_{T_l} - 1)e^{-2\mu(T-T_{l+1})} + e^{-\mu(T-T_{l+1})}\right)\right) \\ &\quad - \frac{2}{Y_{T_l} - 1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\mu(T-T_{l+1})} + \frac{\mathbb{1}_{\{Y_{T_l}>1\}}}{(Y_{T_l} - 1)^2}\end{aligned}$$

$$
\begin{aligned}
&\leq \frac{1}{(Y_{T_l}-1)^2}\mathbb{1}_{\{Y_{T_l}>1\}}\Big(Y_{T_l}e^{-\mu(T-T_{l+1})} + Y_{T_l}(Y_{T_l}-1)e^{-2\mu(T-T_{l+1})}\Big) - \frac{2}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\mu(T-T_{l+1})} \\
&\quad + \frac{\mathbb{1}_{\{Y_{T_l}>1\}}}{(Y_{T_l}-1)^2} \\
&= \frac{\mathbb{1}_{\{Y_{T_l}>1\}}}{Y_{T_l}-1}e^{-\mu(T-T_{l+1})} + \frac{\mathbb{1}_{\{Y_{T_l}>1\}}}{(Y_{T_l}-1)^2}e^{-\mu(T-T_{l+1})} + \frac{Y_{T_l}}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-2\mu(T-T_{l+1})} \\
&\quad - \frac{2}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\mu(T-T_{l+1})} + \frac{\mathbb{1}_{\{Y_{T_l}>1\}}}{(Y_{T_l}-1)^2} \\
&\leq \mathbb{1}_{\{Y_{T_l}>1\}}e^{-2\mu(T-T_{l+1})} + \frac{1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-2\mu(T-T_{l+1})} + \frac{1}{(Y_{T_l}-1)^2}\mathbb{1}_{\{Y_{T_l}>1\}} \\
&\leq \left(\frac{1}{Y_{T_l}-1} + e^{-2\mu(T-T_{l+1})}\right)\mathbb{1}_{\{Y_{T_l}>1\}}.
\end{aligned} \tag{55}
$$

By (54) and (55), we can bound the right-hand side of (53) from above by

$$
\begin{aligned}
\left(\frac{3}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} - \mathbb{1}_{\{Y_{T_l}>1\}}e^{-2\mu(T-T_{l+1})} + e^{-2\mu(T-T_{l+1})}\right)^{\frac{1}{2}} &\leq \left(\frac{3}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} + \mathbb{1}_{\{Y_{T_l}=1\}}\right)^{\frac{1}{2}} \\
&\leq \left(\frac{6}{Y_{T_l}}\right)^{\frac{1}{2}}.
\end{aligned} \tag{56}
$$

☐

The following purely analytical lemma is also used in the proof of Theorem 5.4.

5.18 Lemma
For all $x \geq 0$, we have

$$
\sum_{k=2}^{\infty} \frac{(-x)^k}{k!} \leq \frac{x^2}{2}. \tag{57}
$$

Proof: Let $x \geq 0$. The left-hand side of (57) is equal to $e^{-x} - 1 + x$. For $x = 0$, this expression is equal to $\frac{x^2}{2}$. By $1 - e^{-x} \leq x$, the derivative of the left-hand side of (57) is everywhere smaller than or equal to the derivative of the right-hand side of (57). Hence the statement follows. ☐

For the conditional expectation of the sum of the squared inter-event times since the birth of the randomly picked node, we have the following lemma.

5.19 Lemma
For $T \geq (\frac{2}{\lambda-\mu}\log(2) \vee \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu})$, we have

$$
\mathbb{E}\Bigg(\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(T_{l+1}-T_l)^2 \,\Bigg|\, Y_T > 0\Bigg) \leq \frac{\mu}{\lambda}\frac{T^2}{4}e^{-(\lambda-\mu)T} + 60T^2\frac{\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}e^{-(\lambda+\mu)T} + T^2e^{-\frac{1}{2}\lambda T} \tag{58}
$$

$$
\begin{aligned}
&+ \frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)T\right)T^2e^{-(\lambda-\mu)T} \\
&+ \left(\frac{3}{4}T^2 + \frac{T}{2(\lambda+\mu)}\right)\left(1 + 2\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)T\right)e^{-\frac{1}{4}(\lambda-\mu)T}.
\end{aligned} \tag{59}
$$

Proof: For the left-hand side of (59), we deduce

$$\begin{aligned}
&\mathbb{E}\Big(\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(T_{l+1}-T_l)^2\;\Big|\;Y_T>0\Big)\\
&\leq \mathbb{E}\bigg(\mathbb{1}_{\{T_{\kappa(T)}>T\}}\mathbb{1}_{\{T_{\mathcal{K}(T)}<T_{r(J_T)}\}}\sum_{l=1}^{\mathcal{M}_T-1}\mathbb{1}_{\{T_l\geq T_{r(J_T)}\}}(T_{l+1}-T_l)\max_{r(J_T)\leq j\leq\mathcal{M}_T-1}(T_{j+1}-T_j)\;\bigg|\;Y_T>0\bigg)\\
&\quad+\mathbb{E}(\mathbb{1}_{\{T_{\kappa(T)}\leq T\}}|Y_T>0)T^2+\mathbb{E}(\mathbb{1}_{T_{\mathcal{K}(T)}\geq T_{r(J_T)}\}}|Y_T>0)T^2. \qquad (60)
\end{aligned}$$

Note that, given $(Y_{T_k})_{k\in\mathbb{N}}$ and $\mathcal{K}(T)$, the inter-event times $T_{j+1}-T_j$ are $\mathrm{Exp}((\lambda+\mu)Y_{T_j})$ distributed and independent for $j>\mathcal{K}(T)$. In order to derive an upper bound for the first conditional expectation on the right-hand side of (60), we introduce a sequence of random variables $(U_j)_{j\in\mathbb{N}}$ such that, given $(Y_{T_k})_{k\in\mathbb{N}}$ and $\mathcal{K}(T)$, $U_j\sim\mathrm{Exp}((\lambda+\mu)\min_{\mathcal{K}(T)<k}Y_{T_k})$ i.i.d. Then, given $(Y_{T_k})_{k\in\mathbb{N}}$ and $\mathcal{K}(T)$, we have $T_{j+1}-T_j\leq_{st}U_j$ for $\mathcal{K}(T)<j\leq\kappa(T)$ and obtain

$$\begin{aligned}
&\mathbb{E}\bigg(\mathbb{1}_{\{T_{\kappa(T)}>T\}}\mathbb{1}_{\{T_{\mathcal{K}(T)}<T_{r(J_T)}\}}\sum_{l=1}^{\mathcal{M}_T-1}\mathbb{1}_{\{T_l\geq T_{r(J_T)}\}}(T_{l+1}-T_l)\max_{r(J_T)\leq j\leq\mathcal{M}_T-1}(T_{j+1}-T_j)\;\bigg|\;Y_T>0\bigg)\\
&\leq\frac{T}{2}\mathbb{E}\bigg(\mathbb{1}_{\{T_{\kappa(T)}>T\}}\mathbb{1}_{\{T_{\mathcal{K}(T)}<T_{r(J_T)}\}}\max_{r(J_T)\leq j\leq\mathcal{M}_T-1}(T_{j+1}-T_j)\;\bigg|\;Y_\infty>0\bigg)+\frac{T^2}{4}\mathbb{P}(Y_\infty=0|Y_T>0). \qquad (61)
\end{aligned}$$

By Lemma 5.5, the second summand is equal to

$$\frac{T^2}{4}\frac{\mu}{\lambda}e^{-(\lambda-\mu)T}.$$

The first summand of (61) is bounded from above by

$$\begin{aligned}
&\frac{T}{2}\mathbb{E}\bigg(\mathbb{E}\bigg(\mathbb{1}_{\{T_{\kappa(T)}>T\}}\mathbb{1}_{\{T_{\mathcal{K}(T)}<T_{r(J_T)}\}}\max_{\mathcal{K}(T)<j\leq\kappa(T)}(T_{j+1}-T_j)\;\bigg|\;\mathcal{K}(T),(Y_{T_k})_{k\geq1}\bigg)\;\bigg|\;Y_\infty>0\bigg)\\
&\leq\frac{T}{2}\mathbb{E}\bigg(\mathbb{E}\bigg(\max_{1\leq j\leq\kappa(T)}U_j\;\bigg|\;\mathcal{K}(T),(Y_{T_k})_{k\geq1}\bigg)\;\bigg|\;Y_\infty>0\bigg)\\
&=\frac{T}{2}\mathbb{E}\bigg(\frac{1}{(\lambda+\mu)\min\limits_{\mathcal{K}(T)<k}Y_{T_k}}\sum_{l=1}^{\kappa(T)}\frac{1}{l}\;\bigg|\;Y_\infty>0\bigg),
\end{aligned}$$

where the last equality follows from from the formula for the expectation of the maximum of i.i.d. exponentially distributed random variables (see e.g. the introduction of [Eis08]).

Using the well-known upper bound for the harmonic sum yields

$$\begin{aligned}
&\frac{T}{2}\mathbb{E}\bigg(\frac{1}{(\lambda+\mu)\min\limits_{\mathcal{K}(T)<k}Y_{T_k}}\sum_{l=1}^{\kappa(T)}\frac{1}{l}\;\bigg|\;Y_\infty>0\bigg)\\
&\leq\frac{T}{2(\lambda+\mu)}\mathbb{E}\bigg(\frac{1}{\min\limits_{\mathcal{K}(T)<k}Y_{T_k}}\;\bigg|\;Y_\infty>0\bigg)(\log(\kappa(T))+1)\\
&\leq\frac{T}{2(\lambda+\mu)}\mathbb{E}\bigg(\frac{1}{\min\limits_{\mathcal{K}(T)<k}Y_{T_k}}\;\bigg|\;Y_\infty>0\bigg)\Big(\frac{3}{2}(\lambda+\mu)T+1\Big). \qquad (62)
\end{aligned}$$

For the conditional expectation in (62), we obtain for $T \geq \frac{2\log(2)}{\lambda-\mu}$

$$\begin{aligned}\mathbb{E}\bigg(\frac{1}{\min\limits_{\mathcal{K}(T)<k} Y_{T_k}}\;\bigg|\; Y_\infty > 0\bigg) &= \mathbb{E}\bigg(\frac{1}{\min\limits_{\frac{T}{2}\leq t} Y_t}\;\bigg|\; Y_\infty > 0\bigg)\\ &\leq \bigg(1 + 2\log\bigg(\frac{\lambda}{\lambda-\mu}\bigg) + (\lambda-\mu)T\bigg)e^{-\frac{1}{4}(\lambda-\mu)T}. \end{aligned} \tag{63}$$

where the last line follows from Lemma 5.13 with $\delta = 1$ and $\gamma = \frac{1}{4}$ and Lemma 5.14.

Thus we can conclude that the first summand of the right-hand side of (61) is smaller than or equal to

$$\bigg(\frac{3}{4}T^2 + \frac{T}{2(\lambda+\mu)}\bigg)\bigg(1 + 2\log\bigg(\frac{\lambda}{\lambda-\mu}\bigg) + (\lambda-\mu)T\bigg)e^{-\frac{1}{4}(\lambda-\mu)T}$$

for sufficiently large T.

For the last line of (60), we can use the upper bounds from Lemma 5.10 and obtain that for $T \geq (\frac{1}{\lambda-\mu}\log(2) \vee \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu})$, it is smaller than or equal to

$$\frac{60\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}T^2e^{-(\lambda+\mu)T} + T^2e^{-\frac{1}{2}\lambda T} + \frac{2(\lambda-\mu)}{\lambda}\bigg(\log\bigg(\frac{\lambda}{\lambda-\mu}\bigg) + (\lambda-\mu)T\bigg)T^2e^{-(\lambda-\mu)T}. \tag{64}$$

Altogether, we obtain (59) for $T \geq (\frac{2}{\lambda-\mu}\log(2) \vee \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu})$. ☐

5.2.4 Proof of the main theorem

In order to simplify notation, we introduce $\mathbb{E}^*(\,\cdot\,) = \mathbb{E}(\,\cdot\,|Y_T > 0)$ and $\mathbb{P}^*(\,\cdot\,) = \mathbb{P}(\,\cdot\,|Y_T > 0)$.

In the following we prove Theorem 5.4. As mentioned before, we use Theorem 2.1 to obtain

$$d_{TV}(\mathrm{MixPo}(\Lambda_T^*), \mathrm{MixPo}(\mathrm{M}^*)) \leq \mathbb{E}^*|\Lambda_T - \mathrm{M}|.$$

In order to find an upper bound for $\mathbb{E}^*|\Lambda_T - \mathrm{M}|$, we use the triangle inequality for the absolute value after plugging in the definitions of Λ_T and M given by (27) and (35), which yields

$$\begin{aligned}\mathbb{E}^*\Bigg|&\frac{\alpha(1-e^{-\beta(T-T_{\mathcal{M}_T})})}{(Y_T-1)\beta}\mathbb{1}_{\{Y_T>1\}}\sum_{\substack{i=1\\ i\neq J_T}}^{B_T}(S_i + S_{J_T})\mathbb{1}_{\{T_i^->T\}}\\ &+\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{B_T}(S_i + S_{J_T})\mathbb{1}_{\{T_i^->T\}}\sum_{l=r(i)\vee r(J_T)}^{\mathcal{M}_T-1}(e^{-\beta(T-T_{l+1})} - e^{-\beta(T-T_l)})\frac{1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}\\ &-\frac{\alpha}{\beta+\mu}(\mathbb{E}(S) + S_{J_\infty})(1 - e^{-(\beta+\mu)A_{J_\infty}})\Bigg|.\end{aligned}$$

This is bounded from above by

$$\mathbb{E}^*\bigg(\frac{\alpha(1-e^{-\beta(T-T_{\mathcal{M}_T})})}{(Y_T-1)\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{B_T} S_i\mathbb{1}_{\{T_i^->T\}}\mathbb{1}_{\{Y_T>1\}}\bigg) \tag{65}$$

$$+\,\mathbb{E}^*\bigg(\frac{\alpha(1-e^{-\beta(T-T_{\mathcal{M}_T})})}{(Y_T-1)\beta}S_{J_T}\sum_{\substack{i=1\\ i\neq J_T}}^{B_T}\mathbb{1}_{\{T_i^->T\}}\mathbb{1}_{\{Y_T>1\}}\bigg) \tag{66}$$

$$+\,\mathbb{E}^*\bigg|\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{B_T}(S_i-\mathbb{E}(S))\mathbb{1}_{\{T_i^->T\}}\sum_{l=r(i)\vee r(J_T)}^{\mathcal{M}_T-1}(e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)})\frac{1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}\bigg| \tag{67}$$

$$\begin{aligned}&+\,\mathbb{E}^*\bigg|\frac{\alpha}{\beta}\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\frac{1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}\sum_{\substack{i=1\\ i\neq J_T}}^{r^{-1}(l)}(\mathbb{E}(S)+S_{J_T})\mathbb{1}_{\{T_i^->T\}}(e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)})\\ &\quad-\frac{\alpha}{\beta+\mu}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(e^{-(\beta+\mu)(T-T_{l+1})}-e^{-(\beta+\mu)(T-T_l)})\bigg|\end{aligned} \tag{68}$$

$$\begin{aligned}&+\,\mathbb{E}^*\bigg|\frac{\alpha}{\beta+\mu}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(e^{-(\beta+\mu)(T-T_{l+1})}-e^{-(\beta+\mu)(T-T_l)})\\ &\quad-\frac{\alpha}{\beta+\mu}(\mathbb{E}(S)+S_{J_\infty})(1-e^{-(\beta+\mu)A_{J_\infty}})\bigg|,\end{aligned} \tag{69}$$

where $\mathcal{M}_T$ is the number of events up to time T and $r^{-1}(l)$ is the number of births that occur not later than the lth event for all $l\in\{1,\dots,\mathcal{M}_T\}$, i.e. $r^{-1}:\{1,\dots,\mathcal{M}_T\}\to\{1,\dots,B_T\}, l\mapsto\sum_{i=1}^{B_T}\mathbb{1}_{\{r(i)\le l\}}$.

In the following, we deduce upper bounds for (65)-(69).

Upper bound for (65) and (66)

We treat (65) similarly to the corresponding expression in the pure birth case, but condition on B_T, D_T, J_T and the information which nodes survive up to time T, and obtain

$$\mathbb{E}^*\bigg(\frac{\alpha(1-e^{-\beta(T-T_{\mathcal{M}_T})})}{(Y_T-1)\beta}\mathbb{1}_{\{Y_T>1\}}\sum_{\substack{i=1\\ i\neq J_T}}^{B_T}S_i\mathbb{1}_{\{T_i^->T\}}\bigg)\le\frac{\alpha}{\beta}\mathbb{E}(S)\mathbb{E}^*(1-e^{\beta(T-T_{\mathcal{M}_T})}). \tag{70}$$

Since we condition on J_T, the same expression is also obtained for (66).

Lemma 5.11 reveals an upper bound for the conditional expectation on the right-hand sides of (70): For $T\ge\frac{1}{\lambda-\mu}\log(2)$, we have

$$\mathbb{E}^*(1-e^{-\beta(T-T_{\mathcal{M}_T})})\le\frac{2(\lambda-\mu)}{\lambda e^{(\lambda-\mu)T}}+\frac{\beta}{\lambda}\mathbb{E}^*\bigg(\frac{1}{Y_T-1}\mathbb{1}_{\{Y_T>1\}}\bigg).$$

Lemma 5.12 gives us an upper bound for the second summand of the upper bound from Lemma 5.11:

$$\mathbb{E}^*\bigg(\frac{1}{Y_T-1}\mathbb{1}_{\{Y_T>1\}}\bigg)\le\frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)T}-\mu}\bigg(\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg).$$

Altogether, for $T\ge\frac{1}{\lambda-\mu}\log(2)$, we obtain that the expectation $\mathbb{E}^*(1-e^{-\beta(T-T_{\mathcal{M}_T})})$ is smaller than or equal to

$$\frac{2}{\lambda}\bigg(\lambda-\mu+\beta\bigg(\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)\bigg)e^{-(\lambda-\mu)T}. \tag{71}$$

Plugging this expression in the right-hand side of (70) results in the following upper bound for the sum of (65) and (66):

$$\frac{4\alpha\mathbb{E}(S)}{\beta\lambda}\left(\lambda-\mu+\beta\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)\right)e^{-(\lambda-\mu)T}.$$

Upper bound for (67)

Recall that $R_{T_l,T}$ is the number of nodes that are alive at time T_l and survive up to time T. We compute

$$\begin{aligned}
&\mathbb{E}^*\left(\left|\frac{\alpha}{\beta}\sum_{\substack{i=1\\ i\neq J_T}}^{B_T}(S_i-\mathbb{E}(S))\mathbb{1}_{\{T_i^->T\}}\sum_{l=r(i)\vee r(J_T)}^{\mathcal{M}_T-1}(e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)})\frac{1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}\right|\right)\\
&\leq\mathbb{E}^*\left(\left|\frac{\alpha}{\beta}\sum_{\substack{l=r(J_T)\\ R_{T_l,T}\geq 2}}^{\mathcal{M}_T-1}(e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)})\frac{R_{T_l,T}-1}{Y_{T_l}-1}\frac{\mathbb{1}_{\{Y_{T_l}>1\}}}{R_{T_l,T}-1}\sum_{\substack{i=1\\ i\neq J_T}}^{r^{-1}(l)}(S_i-\mathbb{E}(S))\mathbb{1}_{\{T_i^->T\}}\right|\right)\\
&\leq\mathbb{E}^*\left(\frac{\alpha}{\beta}\sum_{\substack{l=r(J_T)\\ R_{T_l,T}\geq 2}}^{\mathcal{M}_T-1}(e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)})\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}\right.\\
&\qquad\left.\cdot\,\mathbb{E}\left(\frac{1}{R_{T_l,T}-1}\left|\sum_{\substack{i=1\\ i\neq J_T}}^{r^{-1}(l)}(S_i-\mathbb{E}(S))\mathbb{1}_{\{T_i^->T\}}\right|\;\middle|\;(Y_t)_{0\leq t\leq T},J_T\right)\right).
\end{aligned}\tag{72}$$

Note in the second line that we can restrict the sum to $R_{T_l,T}\geq 2$ because $R_{T_l,T}\geq 1$ by $l\geq r(J_T)$ and a summand with $R_{T_l,T}=1$ in the first line would be 0 anyway.

Since the sum in the last line of (72) has exactly $R_{T_l,T}-1$ summands of the form $S_i-\mathbb{E}(S)$ and all other summands are zero, the right-hand side of (72) is smaller than or equal to

$$\mathbb{E}^*\left(\left|\frac{\alpha}{\beta}\sum_{\substack{l=r(J_T)\\ R_{T_l,T}\geq 2}}^{\mathcal{M}_T-1}(e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)})\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}\frac{\sigma_S}{\sqrt{R_{T_l,T}-1}}\right|\right).\tag{73}$$

Since $e^{-\beta(T-T_{l+1})}-e^{-\beta(T-T_l)}\leq\beta(T_{l+1}-T_l)$ and $R_{T_l,T}\leq Y_{T_l}$, the expression (73) is smaller than or equal to

$$\begin{aligned}
\mathbb{E}^*\left(\alpha\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\frac{\sigma_S}{\sqrt{Y_{T_l}-1}}\mathbb{1}_{\{Y_{T_l}>1\}}(T_{l+1}-T_l)\right)&\leq\mathbb{E}^*\left(\alpha\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\frac{\sigma_S}{\sqrt{Y_{T_l}-\frac{Y_{T_l}}{2}}}\mathbb{1}_{\{Y_{T_l}>1\}}(T_{l+1}-T_l)\right)\\
&\leq\sqrt{2}\alpha\sigma_S T\mathbb{E}^*\left(\max_{r(J_T)\leq k\leq\mathcal{M}_T-1}\frac{1}{\sqrt{Y_{T_k}}}\right)\\
&\leq\sqrt{2}\alpha\sigma_S T\mathbb{E}^*\left(\mathbb{1}_{\{\mathcal{K}(T)<r(J_T)\}}\max_{r(J_T)\leq k\leq\mathcal{M}_T-1}\frac{1}{\sqrt{Y_{T_k}}}\right)\\
&\quad+\sqrt{2}\alpha\sigma_S T\mathbb{E}^*\left(\mathbb{1}_{\{\mathcal{K}(T)\geq r(J_T)\}}\max_{r(J_T)\leq k\leq\mathcal{M}_T-1}\frac{1}{\sqrt{Y_{T_k}}}\right),
\end{aligned}\tag{74}$$

where $\mathcal{K}(T) = \max\{k : T_k \leq \frac{T}{2}\}$ (see Definition 5.9). We have

$$\begin{aligned}
&\mathbb{E}^*\left(\mathbb{1}_{\{\mathcal{K}(T)<r(J_T)\}} \max_{r(J_T)\leq k\leq \mathcal{M}_T-1} \frac{1}{\sqrt{Y_{T_k}}}\right) \\
&\leq \mathbb{E}\left(\mathbb{1}_{\{\mathcal{K}(T)<r(J_T)\}} \max_{r(J_T)\leq k\leq \mathcal{M}_T-1} \frac{1}{\sqrt{Y_{T_k}}} \;\middle|\; Y_\infty > 0\right)\mathbb{P}^*(Y_\infty > 0) + 1\cdot\mathbb{P}^*(Y_\infty = 0), \qquad (75)
\end{aligned}$$

where $Y_\infty := \lim\limits_{t\to\infty} Y_t \in \{0,\infty\}$ as before.

By Lemma 5.5, we have for the second summand of the right-hand side of (75)

$$\mathbb{P}^*(Y_\infty = 0) = \frac{\mu}{\lambda}e^{-(\lambda-\mu)T}.$$

The first summand of the right-hand side of (75) is smaller than or equal to

$$\mathbb{E}\left(\max_{\mathcal{K}(T)<k} \frac{1}{\sqrt{Y_{T_k}}} \;\middle|\; Y_\infty > 0\right) = \mathbb{E}\left(\max_{\frac{T}{2}\leq t} \frac{1}{\sqrt{Y_t}} \;\middle|\; Y_\infty > 0\right).$$

By combining Lemma 5.13, where $\delta = \frac{1}{2}$ and $\gamma = \frac{1}{6}$, and Lemma 5.14, we obtain for $T \geq \frac{2}{\lambda-\mu}\log(2)$

$$\begin{aligned}
\mathbb{E}\left(\max_{\frac{T}{2}\leq t} \frac{1}{\sqrt{Y_t}} \;\middle|\; Y_\infty > 0\right) &\leq e^{-\frac{1}{6}(\lambda-\mu)T} + \mathbb{E}\left(\frac{1}{Y_{\frac{T}{2}}} \;\middle|\; Y_\infty > 0\right)e^{\frac{1}{3}(\lambda-\mu)T} \\
&\leq \left(1 + 2\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)T\right)e^{-\frac{1}{6}(\lambda-\mu)T}. \qquad (76)
\end{aligned}$$

Thus for $T \geq \frac{2}{\lambda-\mu}\log(2)$, the first summand of the right-hand side of (74) is smaller than or equal to

$$\sqrt{2}\alpha\frac{\mu}{\lambda}\sigma_S T e^{-(\lambda-\mu)T} + \sqrt{2}\alpha\sigma_S\left(1 + 2\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)T\right)Te^{-\frac{1}{6}(\lambda-\mu)T}. \qquad (77)$$

By Lemma 5.10(i), we obtain that for $T \geq \frac{1}{\lambda-\mu}\log(2)$, the second summand of the right-hand side of (74) is bounded from above by

$$\sqrt{2}\alpha\sigma_S T\left(e^{-\frac{1}{2}\lambda T} + \frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right) + (\lambda-\mu)T\right)e^{-(\lambda-\mu)T}\right). \qquad (78)$$

Thus for $T \geq \frac{2}{\lambda-\mu}\log(2)$, the expression (67) is bounded from above by the sum of (77) and (78).

Upper bound for (68)

For (68), we have

$$\begin{aligned}
&\mathbb{E}^*\Bigg|\frac{\alpha}{\beta}\sum_{l=r(J_T)}^{\mathcal{M}_T-1} \frac{1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} \sum_{\substack{i=1\\ i\neq J_T}}^{r^{-1}(l)} (\mathbb{E}(S) + S_{J_T})\mathbb{1}_{\{T_i^- > T\}}(e^{-\beta(T-T_{l+1})} - e^{-\beta(T-T_l)}) \\
&\qquad - \frac{\alpha}{\beta+\mu}(\mathbb{E}(S) + S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(e^{-(\beta+\mu)(T-T_{l+1})} - e^{-(\beta+\mu)(T-T_l)})\Bigg| \\
&= \mathbb{E}^*\Bigg|\frac{\alpha(\beta+\mu)}{\beta(\beta+\mu)}(\mathbb{E}(S) + S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1} \frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}(e^{-\beta(T-T_{l+1})} - e^{-\beta(T-T_l)})
\end{aligned}$$

$$-\frac{\alpha\beta}{\beta(\beta+\mu)}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(e^{-(\beta+\mu)(T-T_{l+1})}-e^{-(\beta+\mu)(T-T_l)})\Bigg| \tag{79}$$

since the second sum on the left-hand side of (79) has exactly $R_{T_l,T}-1$ non-zero summands. The right-hand side of (79) is equal to

$$\begin{aligned}\mathbb{E}^*\Bigg|\frac{\alpha}{\beta(\beta+\mu)}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\Bigg(&(\beta+\mu)\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\beta(T-T_{l+1})}(1-e^{-\beta(T_{l+1}-T_l)})\\ &-\beta e^{-(\beta+\mu)(T-T_{l+1})}(1-e^{-(\beta+\mu)(T_{l+1}-T_l)})\Bigg)\Bigg|.\end{aligned} \tag{80}$$

Now we use

$$1-e^{-\beta(T_{l+1}-T_l)}=\beta(T_{l+1}-T_l)-\sum_{k=2}^{\infty}\frac{(-\beta(T_{l+1}-T_l))^k}{k!}$$

and

$$1-e^{-(\beta+\mu)(T_{l+1}-T_l)}=(\beta+\mu)(T_{l+1}-T_l)-\sum_{k=2}^{\infty}\frac{(-(\beta+\mu)(T_{l+1}-T_l))^k}{k!}$$

to obtain that (80) is smaller than or equal to

$$\begin{aligned}&\mathbb{E}^*\Bigg|\frac{\alpha}{\beta(\beta+\mu)}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(\beta+\mu)\beta\Bigg(\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}-e^{-\mu(T-T_{l+1})}\Bigg)e^{-\beta(T-T_{l+1})}(T_{l+1}-T_l)\Bigg|\\
&+\mathbb{E}^*\Bigg|\frac{\alpha}{\beta(\beta+\mu)}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\Bigg(\beta e^{-(\beta+\mu)(T-T_{l+1})}\sum_{k=2}^{\infty}\frac{(-(\beta+\mu)(T_{l+1}-T_l))^k}{k!}\\
&\qquad\qquad\qquad\qquad-(\beta+\mu)\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}e^{-\beta(T-T_{l+1})}\sum_{k=2}^{\infty}\frac{(-\beta(T_{l+1}-T_l))^k}{k!}\Bigg)\Bigg|\\
&\leq\mathbb{E}^*\Bigg(\alpha(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\infty}\mathbb{1}_{\{T_{l+1}\leq T\}}\Bigg|\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}-e^{-\mu(T-T_{l+1})}\Bigg|(T_{l+1}-T_l)\Bigg)\\
&\quad+\mathbb{E}^*\Bigg(\frac{\alpha}{\beta+\mu}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\sum_{k=2}^{\infty}\frac{(-(\beta+\mu)(T_{l+1}-T_l))^k}{k!}\Bigg)\\
&\quad+\mathbb{E}^*\Bigg(\frac{\alpha}{\beta}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\sum_{k=2}^{\infty}\frac{(-\beta(T_{l+1}-T_l))^k}{k!}\Bigg)\Bigg).\end{aligned} \tag{81}$$

Firstly, we consider the first summand of the right-hand side of (81). Since the social index S_{J_T} is independent of all other random variables appearing in (81), this summand is equal to

$$2\alpha\mathbb{E}(S)\mathbb{E}^*\Bigg(\sum_{l=r(J_T)}^{\infty}\mathbb{1}_{\{T_{l+1}\leq T\}}\Bigg|\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}-e^{-\mu(T-T_{l+1})}\Bigg|(T_{l+1}-T_l)\Bigg).$$

We derive

$$\mathbb{E}^*\Bigg(\sum_{l=r(J_T)}^{\infty}\mathbb{1}_{\{T_{l+1}<T\}}\Bigg|\frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}}-e^{-\mu(T-T_{l+1})}\Bigg|(T_{l+1}-T_l)\Bigg)$$

$$\begin{aligned}
&\leq \mathbb{E}^*\Bigg(\sum_{l=\mathcal{K}(T)+1}^{\infty} \mathbb{1}_{\{T_{l+1}<T\}} \left| \frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})} \right| (T_{l+1}-T_l) \Bigg) \\
&\quad + T\mathbb{P}^*(\mathcal{K}(T) \geq r(J_T)). \qquad (82)
\end{aligned}$$

For $\frac{1}{\lambda-\mu}\log(2)$, the second summand of the right-hand side of (82) is smaller than or equal to

$$Te^{-\frac{1}{2}\lambda T} + \frac{2(\lambda-\mu)}{\lambda}\bigg(\log\Big(\frac{\lambda}{\lambda-\mu}\Big) + (\lambda-\mu)T\bigg)Te^{-(\lambda-\mu)T}$$

by Lemma 5.10(i). The first summand of the right-hand side of (82) is equal to

$$\mathbb{E}^*\Bigg(\sum_{l=\mathcal{K}(T)+1}^{\infty} \mathbb{E}\bigg(\mathbb{1}_{\{T_{l+1}<T\}} \left| \frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})} \right| (T_{l+1}-T_l) \ \bigg|\ \mathcal{K}(T), Y_T>0 \bigg)\Bigg).$$

For the inner expectation, we have

$$\begin{aligned}
&\mathbb{E}\bigg(\mathbb{1}_{\{T_{l+1}<T\}} \left| \frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})} \right| (T_{l+1}-T_l) \ \bigg|\ \mathcal{K}(T), Y_T>0 \bigg) \\
&= \mathbb{E}\bigg(\mathbb{1}_{\{T_{l+1}<T\}} \left| \frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})} \right| (T_{l+1}-T_l) \ \bigg|\ \mathcal{K}(T), Y_{T_l}>0 \bigg) \frac{\mathbb{P}(Y_{T_l}>0|\mathcal{K}(T))}{\mathbb{P}(Y_T>0|\mathcal{K}(T))}. \qquad (83)
\end{aligned}$$

For the fraction, we obtain for $l \geq \mathcal{K}(T)+1$

$$\frac{\mathbb{P}(Y_{T_l}>0|\mathcal{K}(T))}{\mathbb{P}(Y_T>0|\mathcal{K}(T))} \leq \frac{\mathbb{P}(Y_{T_{\mathcal{K}(T)}}>0|\mathcal{K}(T))}{\mathbb{E}(\mathbb{P}(Y_T>0|\mathcal{K}(T), Y_{T_{\mathcal{K}(T)}})|\mathcal{K}(T))}. \qquad (84)$$

Note that, given $\mathcal{K}(T)$ and $Y_{T_{\mathcal{K}(T)}}$, we know that almost surely $Y_{T_{\mathcal{K}(T)}}$ nodes are alive at time $\frac{T}{2}$ since a jump at this time has probability zero. Thus due to the Markov property for $(Y_t)_{t\geq 0}$ and the formula for the extinction probability of a linear birth and death process with a general initial value given in Remark 3.1, the conditional probability $\mathbb{P}(Y_T>0|\mathcal{K}(T), Y_{T_{\mathcal{K}(T)}})$ is equal to $p_0(\frac{T}{2})^{Y_{T_{\mathcal{K}(T)}}}$, which implies that (84) is equal to

$$\frac{\mathbb{P}(Y_{T_{\mathcal{K}(T)}}>0|\mathcal{K}(T))}{\mathbb{E}(1-p_0(\frac{1}{2}T)^{Y_{T_{\mathcal{K}(T)}}}|\mathcal{K}(T))} \leq \frac{\mathbb{P}(Y_{T_{\mathcal{K}(T)}}>0|\mathcal{K}(T))}{(1-p_0(\frac{1}{2}T))\mathbb{P}(Y_{T_{\mathcal{K}(T)}}>0|\mathcal{K}(T))} = \frac{\lambda e^{\frac{1}{2}(\lambda-\mu)T}-\mu}{(\lambda-\mu)e^{\frac{1}{2}(\lambda-\mu)T}} \leq \frac{\lambda}{\lambda-\mu}.$$

Thus the right-hand side of (83) is smaller than or equal to

$$\mathbb{E}\Bigg(\mathbb{E}\bigg(\mathbb{1}_{\{T_{l+1}<T\}} \left| \frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})} \right| (T_{l+1}-T_l) \ \bigg|\ Y_{T_l}, T_l, \mathcal{K}(T) \bigg) \ \bigg|\ \mathcal{K}(T), Y_{T_l}>0 \Bigg)\frac{\lambda}{\lambda-\mu}. \qquad (85)$$

By the Markov property of $(Y_t)_{t\geq 0}$, we may omit the conditioning on $\mathcal{K}(T)$ if we condition on Y_{T_l} and T_l for $l \geq \mathcal{K}(T)+1$. Conditioning in addition on T_{l+1}, we see that (85) is equal to

$$\mathbb{E}\Bigg(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)\mathbb{E}\bigg(\left| \frac{R_{T_l,T}-1}{Y_{T_l}-1}\mathbb{1}_{\{Y_{T_l}>1\}} - e^{-\mu(T-T_{l+1})} \right| \ \bigg|\ Y_{T_l}, T_l, T_{l+1} \bigg) \ \bigg|\ \mathcal{K}(T), Y_{T_l}>0 \Bigg)\frac{\lambda}{\lambda-\mu}. \qquad (86)$$

By Lemma 5.17, summing over l and taking the expectation yields the following upper bound for the first summand of the right-hand side of (82):

$$\mathbb{E}^*\bigg(\sum_{l=\mathcal{K}(T)+1}^{\infty}\sqrt{6}\frac{\lambda}{\lambda-\mu}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T),Y_{T_l}>0\Big)\bigg). \tag{87}$$

Obviously, (87) is equal to

$$\begin{aligned}&\sqrt{6}\frac{\lambda}{\lambda-\mu}\mathbb{E}^*\bigg(\mathbb{1}_{\{Y_\infty>0\}}\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T),Y_{T_l}>0\Big)\bigg)\\&+\sqrt{6}\frac{\lambda}{\lambda-\mu}\mathbb{E}^*\bigg(\mathbb{1}_{\{Y_\infty=0\}}\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T),Y_{T_l}>0\Big)\bigg).\end{aligned}$$

We may conclude that for for $T\geq\frac{1}{\lambda-\mu}\log(2)$, the first summand of (81) is bounded from above by

$$\begin{aligned}2\alpha\mathbb{E}(S)\Bigg(&\sqrt{6}\frac{\lambda}{\lambda-\mu}\mathbb{E}^*\bigg(\mathbb{1}_{\{Y_\infty>0\}}\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T),Y_{T_l}>0\Big)\bigg)\\&+\sqrt{6}\frac{\lambda}{\lambda-\mu}\mathbb{E}^*\bigg(\mathbb{1}_{\{Y_\infty=0\}}\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T),Y_{T_l}>0\Big)\bigg)\\&+Te^{-\frac{1}{2}\lambda T}+\frac{2(\lambda-\mu)}{\lambda}\Big(\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\Big)Te^{-(\lambda-\mu)T}\Bigg).\end{aligned} \tag{88}$$

For the first outer expectation in (88), we have

$$\begin{aligned}&\mathbb{E}^*\bigg(\mathbb{1}_{\{Y_\infty>0\}}\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T),Y_{T_l}>0\Big)\bigg)\\ \leq\;&\mathbb{E}\bigg(\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T)\Big)\,\Big|\,Y_\infty>0\bigg)\\ \leq\;&\mathbb{E}\bigg(\max_{\frac{T}{2}\leq t}Y_t^{-\frac{1}{2}}\mathbb{E}\Big(\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)\,\Big|\,\mathcal{K}(T)\Big)\,\Big|\,Y_\infty>0\bigg)\\ \leq\;&\frac{T}{2}\mathbb{E}\Big(\max_{\frac{T}{2}\leq t}Y_t^{-\frac{1}{2}}\,\Big|\,Y_\infty>0\Big).\end{aligned}$$

By inequality (76), we obtain that for $T\geq\frac{2}{\lambda-\mu}\log(2)$, this expression is smaller than or equal to

$$\frac{T}{2}\Big(1+2\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\Big)e^{-\frac{1}{6}(\lambda-\mu)T}.$$

For the second outer expectation in (88), we obtain

$$\begin{aligned}&\mathbb{E}\bigg(\mathbb{1}_{\{Y_\infty=0\}}\sum_{l=\mathcal{K}(T)+1}^{\infty}\mathbb{E}\Big(\mathbb{1}_{\{T_{l+1}<T\}}(T_{l+1}-T_l)Y_{T_l}^{-\frac{1}{2}}\,\Big|\,\mathcal{K}(T),Y_{T_l}>0\Big)\,\Big|\,Y_T>0\bigg)\\&\leq\frac{T}{2}\mathbb{P}(Y_\infty=0|Y_T>0)=\frac{T}{2}\frac{\mu}{\lambda}e^{-(\lambda-\mu)T},\end{aligned}$$

where the last equality follows from Lemma 5.5.

In conclusion, for $T \geq \frac{2}{\lambda-\mu}\log(2)$, the first summand of the right-hand side of (81) is bounded from above by

$$\begin{aligned}&\sqrt{6}\alpha\frac{\lambda}{\lambda-\mu}\mathbb{E}(S)T\Bigg(\bigg(1+2\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)e^{-\frac{1}{6}(\lambda-\mu)T}+e^{-(\lambda-\mu)T}\Bigg)\\&+2\alpha\mathbb{E}(S)\Bigg(Te^{-\frac{1}{2}\lambda T}+\frac{2(\lambda-\mu)}{\lambda}\bigg(\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)Te^{-(\lambda-\mu)T}\Bigg).\end{aligned}$$

Since the social index S_{J_T} is independent of all other random variables appearing in (81), the second summand of the right-hand side of (81) is smaller than or equal to

$$\frac{2\alpha}{\beta+\mu}\mathbb{E}(S)\mathbb{E}^*\Big(\sum_{l=r(J_T)}^{\mathcal{M}_T-1}\sum_{k=2}^{\infty}\frac{(-(\beta+\mu)(T_{l+1}-T_l))^k}{k!}\Big).$$

By Lemma 5.18, this expression is bounded from above by

$$\frac{\alpha}{\beta+\mu}\mathbb{E}(S)\mathbb{E}^*\Big(\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(\beta+\mu)^2(T_{l+1}-T_l)^2\Big)\Big). \tag{89}$$

For $T \geq (\frac{2}{\lambda-\mu}\log(2) \vee \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu})$, Lemma 5.19 implies that (89), which is an upper bound for the second summand of the right-hand side of (81), is smaller than or equal to

$$\begin{aligned}\alpha(\beta+\mu)\mathbb{E}(S)\Bigg(&\frac{\mu}{\lambda}\frac{T^2}{4}e^{-(\lambda-\mu)T}+\frac{60\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}T^2e^{-(\lambda+\mu)T}+T^2e^{-\frac{1}{2}\lambda T}\\&+\frac{2(\lambda-\mu)}{\lambda}\bigg(\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)T^2e^{-(\lambda-\mu)T}\\&+\bigg(\frac{3}{4}T^2+\frac{T}{2(\lambda+\mu)}\bigg)\bigg(1+2\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)e^{-\frac{1}{4}(\lambda-\mu)T}\Bigg).\end{aligned}$$

For the third summand of the right-hand side of (81), we obviously obtain the same upper bound, but may replace the factor $\beta+\mu$ by β.

We may conclude that for $T \geq (\frac{2}{\lambda-\mu}\log(2) \vee \frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu})$, the expression (68) is bounded from above by

$$\begin{aligned}&\sqrt{6}\alpha\frac{\lambda}{\lambda-\mu}\mathbb{E}(S)T\Bigg(\bigg(1+2\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)e^{-\frac{1}{6}(\lambda-\mu)T}+e^{-(\lambda-\mu)T}\Bigg)\\&+2\alpha\mathbb{E}(S)\Bigg(Te^{-\frac{1}{2}\lambda T}+\frac{2(\lambda-\mu)}{\lambda}\bigg(\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)Te^{-(\lambda-\mu)T}\Bigg)\\&+\alpha(2\beta+\mu)\mathbb{E}(S)\Bigg(\frac{\mu}{\lambda}\frac{T^2}{4}e^{-(\lambda-\mu)T}+\frac{60\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}T^2e^{-(\lambda+\mu)T}+T^2e^{-\frac{1}{2}\lambda T}\\&\qquad+\frac{2(\lambda-\mu)}{\lambda}\bigg(\log\Big(\frac{\lambda}{\lambda-\mu}\Big)+(\lambda-\mu)T\bigg)T^2e^{-(\lambda-\mu)T}\end{aligned}$$

$$+\left(\frac{3}{4}T^2+\frac{T}{2(\lambda+\mu)}\right)\left(1+2\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)e^{-\frac{1}{4}(\lambda-\mu)T}\Bigg).$$

Upper bound for (69)

Recall that we arranged $S_{J_T}=S_{J_\infty}$. Since the sum in (69) telescopes, this implies

$$\begin{aligned}
&\mathbb{E}^*\Bigg(\Bigg|\frac{\alpha}{\beta+\mu}(\mathbb{E}(S)+S_{J_T})\sum_{l=r(J_T)}^{\mathcal{M}_T-1}(e^{-(\beta+\mu)(T-T_{l+1})}-e^{-(\beta+\mu)(T-T_l)})\\
&\qquad-\frac{\alpha}{\beta+\mu}(\mathbb{E}(S)+S_{J_\infty})(1-e^{-(\beta+\mu)A_{J_\infty}})\Bigg|\Bigg)\\
&=\mathbb{E}^*\left(\left|\frac{\alpha}{\beta+\mu}(\mathbb{E}(S)+S_{J_\infty})\left(e^{-(\beta+\mu)(T-T_{\mathcal{M}_T})}-e^{-(\beta+\mu)(T-T_{r(J_T)})}-(1-e^{-(\beta+\mu)A_{J_\infty}})\right)\right|\right)\\
&\le\frac{2\alpha}{\beta+\mu}\mathbb{E}(S)\left(\mathbb{E}^*\left(1-e^{-(\beta+\mu)(T-T_{\mathcal{M}_T})}\right)+\mathbb{E}^*\left(\left|e^{-(\beta+\mu)A_{J_T}(T)}-e^{-(\beta+\mu)A_{J_\infty}}\right|\right)\right),
\end{aligned}\tag{90}$$

where the last inequality holds since S_{J_∞} is independent of all other random variables appearing in (90).

By combining Lemma 5.11 and Lemma 5.12 as before, we obtain for the first conditional expectation on the right-hand side of (90) the upper bound

$$\frac{2}{\lambda}\left(\lambda-\mu+(\beta+\mu)\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)\right)e^{-(\lambda-\mu)T}$$

if $T\ge\frac{1}{\lambda-\mu}\log(2)$.

For the second conditional expectation on the right-hand side of (90), Corollary 3.14 implies

$$\mathbb{E}^*\left(\left|e^{-(\beta+\mu)A_{J_T}(T)}-e^{-(\beta+\mu)A_{J_\infty}}\right|\right)\le\left(\frac{2\lambda}{\beta+\mu+\lambda}+\frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)\right)e^{-(\lambda-\mu)T}$$

if $T\ge\frac{1}{\lambda-\mu}\log(2)$.

Altogether, we obtain that the right-hand side of (90) is smaller than or equal to

$$\frac{4\alpha}{\beta+\mu}\mathbb{E}(S)\left(\frac{\lambda-\mu}{\lambda}+\frac{\beta+\lambda}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)+\frac{\lambda}{\beta+\mu+\lambda}\right)e^{-(\lambda-\mu)T}$$

if $T\ge\frac{1}{\lambda-\mu}\log(2)$.

Conclusion

Combining the upper bounds obtained for (65)-(69), we have for $T\ge(\frac{2}{\lambda-\mu}\log(2)\vee\frac{2(\log(4\lambda)-\log(\lambda-\mu))}{\lambda+\mu})$

$$\begin{aligned}
&d_{TV}(\mathrm{MixPo}(\Lambda_T^*),\mathrm{MixPo}(\mathrm{M}^*))\\
&\le\frac{4\alpha\mathbb{E}(S)}{\beta\lambda}\left(\lambda-\mu+\beta\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)\right)e^{-(\lambda-\mu)T}\\
&+\sqrt{2}\alpha\frac{\mu}{\lambda}\sigma_S Te^{-(\lambda-\mu)T}+\sqrt{2}\alpha\sigma_S\left(1+2\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)Te^{-\frac{1}{6}(\lambda-\mu)T}
\end{aligned}$$

$$
\begin{aligned}
&+\sqrt{2}\alpha\sigma_S T\left(e^{-\frac{1}{2}\lambda T}+\frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)e^{-(\lambda-\mu)T}\right)\\
&\sqrt{6}\alpha\frac{\lambda}{\lambda-\mu}\mathbb{E}(S)T\left(\left(1+2\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)e^{-\frac{1}{6}(\lambda-\mu)T}+e^{-(\lambda-\mu)T}\right)\\
&+2\alpha\mathbb{E}(S)\left(Te^{-\frac{1}{2}\lambda T}+\frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)Te^{-(\lambda-\mu)T}\right)\\
&+\alpha(2\beta+\mu)\mathbb{E}(S)\left(\left(\frac{3}{4}T^2+\frac{T}{2(\lambda+\mu)}\right)\left(1+2\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)e^{-\frac{1}{4}(\lambda-\mu)T}+\frac{T^2}{4}\frac{\mu}{\lambda}e^{-(\lambda-\mu)T}\right)\\
&+\alpha(2\beta+\mu)\mathbb{E}(S)T^2\left(\frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)e^{-(\lambda-\mu)T}+\frac{60\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}e^{-(\lambda+\mu)T}+e^{-\frac{1}{2}\lambda T}\right)\\
&+\frac{4\alpha}{\beta+\mu}\mathbb{E}(S)\left(\frac{\lambda-\mu}{\lambda}+\frac{\beta+\lambda}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)+\frac{\lambda}{\beta+\mu+\lambda}\right)e^{-(\lambda-\mu)T}\\
&=\left(\sqrt{2}\alpha\sigma_S+\sqrt{6}\alpha\frac{\lambda}{\lambda-\mu}\mathbb{E}(S)\right)\left(1+2\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)Te^{-\frac{1}{6}(\lambda-\mu)T}\\
&\quad+\frac{\alpha}{2}(2\beta+\mu)\mathbb{E}(S)\left(\frac{3}{2}T+\frac{1}{\lambda+\mu}\right)\left(1+2\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)Te^{-\frac{1}{4}(\lambda-\mu)T}\\
&\quad+\left(\sqrt{2}\alpha\sigma_S+2\alpha\mathbb{E}(S)+\alpha(2\beta+\mu)\mathbb{E}(S)T\right)Te^{-\frac{1}{2}\lambda T}\\
&\quad+\left(\alpha\mathbb{E}(S)\left(\frac{4}{\beta\lambda}\left(\lambda-\mu+\beta\left(1+\frac{\beta+\lambda}{\beta+\mu}+\left(1+\frac{2\beta+\mu}{2}T\right)(\lambda-\mu)T\right)\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)\right)\right.\right.\\
&\qquad\qquad\left.+\sqrt{6}\frac{\lambda}{\lambda-\mu}T+\frac{(2\beta+\mu)\mu}{4\lambda}T^2+\frac{4(\lambda-\mu)}{\lambda(\beta+\mu)}+\frac{4}{\beta+\mu}\frac{\lambda}{\beta+\lambda+\mu}\right)\\
&\qquad\left.+\sqrt{2}\alpha\sigma_S\left(\frac{\mu}{\lambda}+\frac{2(\lambda-\mu)}{\lambda}\left(\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T\right)\right)T\right)e^{-(\lambda-\mu)T}\\
&\quad+60\alpha\frac{\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}(2\beta+\mu)\mathbb{E}(S)T^2e^{-(\lambda+\mu)T}
\end{aligned}
\tag{91}
$$

In the remainder of the proof, we find a simpler upper bound by elementary calculations. In order to do so, we assume $T \geq \frac{2\log(4(\lambda(\lambda-\mu)^{-1}))}{\lambda-\mu}$. Then we have

$$
\begin{aligned}
&1+2\log\left(\frac{\lambda}{\lambda-\mu}\right)+(\lambda-\mu)T \leq \frac{5}{2}(\lambda-\mu)T,\\
&e^{-\frac{1}{4}(\lambda-\mu)T}=e^{-\frac{1}{6}(\lambda-\mu)T}e^{-\frac{1}{12}(\lambda-\mu)T}\leq\left(4^{-\frac{1}{6}}\wedge\frac{12}{(\lambda-\mu)T}\right)e^{-\frac{1}{6}(\lambda-\mu)T}\leq\left(\frac{2}{5}\wedge\frac{12}{(\lambda-\mu)T}\right)e^{-\frac{1}{6}(\lambda-\mu)T},\\
&e^{-\frac{1}{2}(\lambda-\mu)T}=e^{-\frac{1}{6}(\lambda-\mu)T}e^{-\frac{1}{3}(\lambda-\mu)T}\leq 4^{-\frac{2}{3}}e^{-\frac{1}{6}(\lambda-\mu)T}\leq\frac{2}{5}e^{-\frac{1}{6}(\lambda-\mu)T}
\end{aligned}
$$

and finally $2 \leq (\lambda-\mu)T$. Thus the first three lines of the right-hand side of (91) are smaller than or equal to

$$
\alpha\left(\left(\frac{5\sqrt{6}}{2}\frac{\lambda}{\lambda-\mu}+\frac{2}{5}+\left(\beta+\frac{\mu}{2}\right)\left(\frac{229}{5(\lambda-\mu)}+\frac{2}{\lambda+\mu}\right)\right)\mathbb{E}(S)+\frac{27}{10}\sqrt{2}\sigma_S\right)(\lambda-\mu)T^2e^{-\frac{1}{6}(\lambda-\mu)T}. \tag{92}
$$

Note that we have the following inequalities:

$$\begin{aligned}
&\log\Big(\frac{\lambda}{\lambda-\mu}\Big) + (\lambda-\mu)T \le \frac{3}{2}(\lambda-\mu)T,\\
&1+\log\Big(\frac{\lambda}{\lambda-\mu}\Big) + (\lambda-\mu)T \le 2(\lambda-\mu)T,\\
&e^{-(\lambda+\mu)T} = e^{-2\mu T}e^{-(\lambda-\mu)T} \le \frac{1}{2\mu}\frac{1}{T}e^{-(\lambda-\mu)T}.
\end{aligned}$$

Thus the last four lines of (91) are smaller than or equal to

$$\begin{aligned}
&\alpha\Bigg(\frac{4(\lambda-\mu)}{\beta\lambda}\Big(1+\frac{3}{2}\beta\Big(1+\frac{\beta+\lambda}{\beta+\mu}+\Big(1+\frac{2\beta+\mu}{2}T\Big)(\lambda-\mu)T\Big)T\Big)\mathbb{E}(S)\\
&\quad+\sqrt{2}\Big(\frac{\mu}{\lambda}T+\frac{3(\lambda-\mu)^2}{\lambda}T^2\Big)\sigma_S+\Big(\frac{\beta}{2}+\frac{\mu}{4}\Big)\frac{\mu}{\lambda}\mathbb{E}(S)T^2+\sqrt{6}\frac{\lambda}{\lambda-\mu}\mathbb{E}(S)T\\
&\quad+\frac{4(\lambda-\mu)}{\lambda(\beta+\mu)}\mathbb{E}(S)+\frac{30\lambda^3(\lambda+\mu)}{\mu(\lambda-\mu)^4}(2\beta+\mu)\mathbb{E}(S)T+\frac{4}{\beta+\mu}\frac{\lambda}{\beta+\lambda+\mu}\mathbb{E}(S)\Bigg)e^{-(\lambda-\mu)T}. \qquad (93)
\end{aligned}$$

We first consider those terms in (93) that depend on $\mathbb{E}(S)$. Since $2\le(\lambda-\mu)T$, grouping the terms where β dominates in the denominator, we have

$$\begin{aligned}
&\frac{8\alpha\mathbb{E}(S)}{\beta}\frac{\lambda-\mu}{\lambda}e^{-(\lambda-\mu)T}+\frac{4\alpha\lambda\mathbb{E}(S)}{\beta(\beta+\lambda+\mu)}e^{-(\lambda-\mu)T}+\frac{6(\lambda-\mu)}{\beta}Te^{-(\lambda-\mu)T}\\
&\le\frac{\alpha\mathbb{E}(S)}{\beta}\Big(\frac{2(\lambda-\mu)^2}{\lambda}+2(\lambda-\mu)+6(\lambda-\mu)\Big)Te^{-(\lambda-\mu)T}\\
&\le\frac{\alpha\mathbb{E}(S)}{\beta}10(\lambda-\mu)Te^{-(\lambda-\mu)T}\\
&\le\frac{5\alpha\mathbb{E}(S)}{\beta}(\lambda-\mu)^2T^2e^{-(\lambda-\mu)T}; \qquad (94)
\end{aligned}$$

grouping the β-free terms in (93) yields

$$\begin{aligned}
&\alpha\mathbb{E}(S)\Big(12\frac{\lambda-\mu}{\lambda}T+3\frac{(\lambda-\mu)^2\mu}{\lambda}T^3+3\frac{(\lambda-\mu)^2}{\lambda}T^2+\frac{\mu^2}{4\lambda}T^2+\sqrt{6}\frac{\lambda}{\lambda-\mu}T+\frac{30\lambda^3(\lambda+\mu)}{(\lambda-\mu)^4}T\Big)e^{-(\lambda-\mu)T}\\
&\le 12\alpha\mathbb{E}(S)\frac{\lambda-\mu}{\lambda}Te^{-(\lambda-\mu)T}+\alpha\mathbb{E}(S)\Big(\frac{3(\lambda-\mu)^2}{\lambda}(\mu T+1)+\frac{\mu^2}{4\lambda}+\frac{\sqrt{6}\lambda}{2}+\frac{15\lambda^3(\lambda+\mu)}{(\lambda-\mu)^3}\Big)T^2e^{-(\lambda-\mu)T}\\
&\le\alpha\mathbb{E}(S)\Big(\frac{3(\lambda-\mu)^2}{\lambda}(\mu T+3)+\frac{\mu^2}{4\lambda}+\frac{\sqrt{6}\lambda}{2}+\frac{15\lambda^3(\lambda+\mu)}{(\lambda-\mu)^3}\Big)T^2e^{-(\lambda-\mu)T}; \qquad (95)
\end{aligned}$$

and considering the terms in (93) where β dominates in the numerator, we obtain

$$\begin{aligned}
&\alpha\beta\mathbb{E}(S)\frac{\mu}{2\lambda}T^2e^{-(\lambda-\mu)T}+16\alpha\beta\mathbb{E}(S)\frac{\lambda^3(\lambda+\mu)}{\mu(\lambda-\mu)^4}Te^{-(\lambda-\mu)T}+6\alpha\beta\mathbb{E}(S)\frac{(\lambda-\mu)^2}{\lambda}T^3e^{-(\lambda-\mu)T}\\
&\le\alpha\beta\mathbb{E}(S)\Big(\frac{\mu}{2\lambda}+\frac{8\lambda^3(\lambda+\mu)}{\mu(\lambda-\mu)^3}+6\alpha\beta\frac{(\lambda-\mu)^2}{\lambda}T\Big)T^2e^{-(\lambda-\mu)T}. \qquad (96)
\end{aligned}$$

For the one term in (93) that depends on σ_S, the inequality $2\le(\lambda-\mu)T$ implies

$$\sqrt{2}\Big(\frac{\mu}{\lambda}T+\frac{3(\lambda-\mu)^2}{\lambda}T^2\Big)\sigma_Se^{-(\lambda-\mu)T}\le\sqrt{2}\Big(\frac{\mu(\lambda-\mu)}{2\lambda}+\frac{3(\lambda-\mu)^2}{\lambda}\Big)\sigma_ST^2e^{-(\lambda-\mu)T}$$

$$\begin{aligned}
&= \sqrt{2}\frac{6\lambda^2 - 5\mu\lambda + \mu^2}{2\lambda}\sigma_S T^2 e^{-(\lambda-\mu)T} \\
&\le 3\sqrt{2}\lambda\sigma_S T^2 e^{-(\lambda-\mu)T}. \qquad (97)
\end{aligned}$$

Combining (94)-(97), we obtain the following upper bound for the right-hand side of (93):

$$\begin{aligned}
&\alpha\beta\Bigg(\Big(\frac{1}{2}+\frac{\mu}{4\beta}\Big)\frac{\mu}{\lambda} + \Big(\frac{8}{\mu}+\frac{4}{\beta}\Big)\frac{\lambda^3(\lambda+\mu)}{(\lambda-\mu)^3} + \Big(6T+\frac{\sqrt{6}}{2}+\frac{3}{\beta}(\mu T+3)+\frac{5\lambda}{\beta^2}\Big)\lambda\Bigg)\mathbb{E}(S)T^2e^{-(\lambda-\mu)T} \\
&+ 3\sqrt{2}\alpha\lambda\sigma_S T^2 e^{-(\lambda-\mu)T}. \qquad (98)
\end{aligned}$$

The total upper bound for the theorem is the sum of (92) and (98) and this sum is obviously of the desired order. ☐

6 Negligibility of multiple edges

In this section, we show that multiple edges are negligible (Lemma 6.1 below) and use this result to prove Corollary 1.4 from the introduction, which states our main result for the case where multiple edges are ignored.

6.1 Lemma

The probability that an individual picked uniformly at random at time T has at least one multiple edge given the number of nodes is positive at time T is of order $O(T^2 e^{-\frac{1}{6}(\lambda-\mu)T})$ as $T \to \infty$.

Proof. Let N_T denote the degree of the randomly picked node J_T at time T, i.e. the number of edges that are incident to the node picked uniformly at random at time T. Given N_T, let $\rho_1 < \ldots < \rho_{N_T}$ be the birth times of these edges. Supposing we condition on J_T, N_T, $\rho_1 < \ldots < \rho_{N_T}$ and $(Y_t)_{0\le t\le T}$ and are at time ρ_i, the probability that we do not create a multiple edge that stays a multiple edge up to time T is

$$\Big(1 - \frac{i-1}{Y_{\rho_i}-1}\Big)\mathbb{1}_{\{Y_{\rho_i}\ge i+1\}}.$$

Thus we obtain for the probability that J_T has at least one multiple edge is

$$\begin{aligned}
&\mathbb{E}\Bigg(1 - \Big(1-\frac{1}{Y_{\rho_2}-1}\Big)\mathbb{1}_{\{Y_{\rho_2}\ge 3\}}\cdot\Big(1-\frac{2}{Y_{\rho_3}-1}\Big)\mathbb{1}_{\{Y_{\rho_3}\ge 4\}}\cdot\ldots\cdot\Big(1-\frac{N_T-1}{Y_{\rho_{N_T}}-1}\Big)\mathbb{1}_{\{Y_{\rho_{N_T}}\ge N_T+1\}}\Bigg| Y_T>0\Bigg) \\
&\le \mathbb{E}\Bigg(1-\Bigg(1-\frac{N_T-1}{\min\limits_{T-A_{J_T}(T)\le t\le T} Y_t - 1}\Bigg)^{N_T-1}\mathbb{1}_{\{\min_{T-A_{J_T}(T)\le t\le T} Y_t - 1\ge N_T\}}\Bigg| Y_T>0\Bigg) \\
&\le \mathbb{P}(N_T \ge e^{\frac{1}{12}(\lambda-\mu)T}|Y_T>0) + \mathbb{P}\Bigg(\min_{T-A_{J_T}(T)\le t\le T} Y_t - 1 \le e^{\frac{1}{3}(\lambda-\mu)T}\Bigg|Y_T>0\Bigg) \\
&\quad + 1 - \Bigg(1-\frac{e^{\frac{1}{12}(\lambda-\mu)T}-1}{e^{\frac{1}{3}(\lambda-\mu)T}}\Bigg)^{e^{\frac{1}{12}(\lambda-\mu)T}-1}. \qquad (99)
\end{aligned}$$

Since the asymptotic degree distribution is MixPo(M) with M given by (35), we obtain by conditioning on M that the second moment $\mathbb{E}(N_T^2)$ is equal to

$$\frac{2\alpha E(S)}{\lambda+\beta+\mu} + \frac{2\alpha\mathbb{E}((S+\mathbb{E}(S))^2)}{(\lambda+\beta+\mu)(\lambda+2(\beta+\mu))} \text{ (cf. Subsection 3.3 in [BL10]).}$$

Thus Theorem 5.4 and the Markov inequality imply

$$\begin{aligned}\mathbb{P}(N_T \geq e^{\frac{1}{12}(\lambda-\mu)T} | Y_T > 0) &\leq \mathbb{E}(N_T^2) T^{\frac{1}{4}} e^{-\frac{1}{8}(\lambda-\mu)T} + O(T^2 e^{-\frac{1}{6}(\lambda-\mu)T}) \\ &\leq \left(\frac{2\alpha E(S)}{\lambda+\beta+\mu} + \frac{2\alpha \mathbb{E}((S+\mathbb{E}(S))^2)}{(\lambda+\beta+\mu)(\lambda+2(\beta+\mu))} \right) e^{-\frac{1}{6}(\lambda-\mu)T} + O(T^2 e^{-\frac{1}{6}(\lambda-\mu)T}) \\ &= O(T^2 e^{-\frac{1}{6}(\lambda-\mu)T}).\end{aligned}$$

For the second summand of the right-hand side of (99), we obtain

$$\begin{aligned}\mathbb{P}\left(\min_{T-A_{J_T}(T) \leq t \leq T} Y_t - 1 \leq e^{\frac{1}{3}(\lambda-\mu)T} \middle| Y_T > 0 \right) &\leq \mathbb{P}\left(\max_{\frac{T}{2} \leq t \leq T} \frac{1}{Y_t} > \frac{1}{e^{\frac{1}{3}(\lambda-\mu)T}+1} \middle| Y_T > 0 \right) \\ &\quad + \mathbb{P}\left(A_{J_T}(T) < \frac{T}{2} \,\middle|\, Y_T > 0 \right). \end{aligned} \tag{100}$$

By Lemma 5.10, the second summand of the right-hand side is of order $O(e^{-\frac{1}{2}(\lambda-\mu)T})$ as $T \to \infty$. With $Y_\infty = \lim_{t\to\infty} Y_t \in \{0, \infty\}$, we have

$$\begin{aligned}\mathbb{P}\left(\max_{\frac{T}{2} \leq t \leq T} \frac{1}{Y_t} > \frac{1}{e^{\frac{1}{3}(\lambda-\mu)T}+1}, Y_\infty > 0 \middle| Y_T > 0 \right) &= \frac{\mathbb{P}\left(\max_{\frac{T}{2} \leq t \leq T} \frac{1}{Y_t} > \frac{1}{e^{\frac{1}{3}(\lambda-\mu)T}+1}, Y_\infty > 0 \right)}{\mathbb{P}(Y_T > 0)} \\ &\leq \frac{\mathbb{P}\left(\max_{\frac{T}{2} \leq t \leq T} \frac{1}{Y_t} > \frac{1}{e^{\frac{1}{3}(\lambda-\mu)T}+1}, Y_\infty > 0 \right)}{\mathbb{P}(Y_\infty > 0)} \\ &\leq \mathbb{P}\left(\max_{\frac{T}{2} \leq t \leq T} \frac{1}{Y_t} > \frac{1}{e^{\frac{1}{3}(\lambda-\mu)T}+1} \middle| Y_\infty > 0 \right).\end{aligned}$$

Thus the first summand of the right-hand side of (100) is smaller than or equal to

$$\mathbb{P}\left(\max_{\frac{T}{2} \leq t \leq T} \frac{1}{Y_t} > \frac{1}{e^{\frac{1}{3}(\lambda-\mu)T}+1} \middle| Y_\infty > 0 \right) + \mathbb{P}(Y_\infty = 0 | Y_T > 0). \tag{101}$$

The second summand is smaller than or equal to $\frac{\mu}{\lambda} e^{-(\lambda-\mu)T}$ by Lemma 5.5. By Corollary 5.8 and the inequality (2) in Theorem 6.14 on page 99 in [Yeh95], the first summand of (101) is bounded from above by

$$\mathbb{E}\left(\frac{1}{Y_{\frac{T}{2}}} \middle| Y_\infty > 0 \right) (e^{\frac{1}{3}(\lambda-\mu)T} + 1) = O(Te^{-\frac{1}{2}(\lambda-\mu)T})(e^{\frac{1}{3}(\lambda-\mu)T} + 1) = O(Te^{-\frac{1}{6}(\lambda-\mu)T}),$$

where the first equality follows from Lemma 5.14.

In order to obtain an upper bound for the third summand on the right-hand side of (99), we apply the following general lemma, which can be proved by elementary calculations; cf. [AE06], proof of Theorem 6.23.

6.2 Lemma

Let $(z_n)_{n\in\mathbb{N}}$ be a non-negative sequence. Then for all $n \in \mathbb{N}$ with $\frac{z_n}{n} \leq 1$, we have

$$\left| e^{-z_n} - \left(1 - \frac{z_n}{n} \right)^n \right| \leq \frac{e}{2} \frac{z_n^2}{n}.$$

The third summand of the right-hand side of (99) is smaller than or equal to

$$1-\left(1-\frac{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil}{e^{\frac{1}{3}(\lambda-\mu)T}}\right)^{\lceil e^{(\lambda-\mu)T/12}-1\rceil}, \tag{102}$$

where $\lceil\cdot\rceil$ denotes the ceiling function. Let $T\geq\frac{12\log(2)}{\lambda-\mu}$. Then (102) is bounded from above by

$$\begin{aligned}&1-\left(1-\frac{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil}{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil^4}\right)^{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil}=1-\left(1-\frac{1}{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil^3}\right)^{\lceil e^{(\lambda-\mu)T/12}-1\rceil}\\&\leq 1-\exp\left(-\frac{1}{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil^2}\right)\\&\quad+\left|\exp\left(-\frac{1}{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil^2}\right)-\left(1-\frac{1}{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil^3}\right)^{\lceil e^{(\lambda-\mu)T/12}-1\rceil}\right|.\end{aligned} \tag{103}$$

Thus by Lemma 6.2, the right-hand side of (103) is bounded from above by

$$\frac{1}{\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil^2}+\frac{e}{2\lceil e^{\frac{1}{12}(\lambda-\mu)T}-1\rceil^3}\leq\frac{4}{e^{\frac{1}{6}(\lambda-\mu)T}}+\frac{4e}{e^{\frac{1}{4}(\lambda-\mu)T}}=O(e^{-\frac{1}{6}(\lambda-\mu)T}).$$

Altogether, we obtain that the probability that J_T has at least one multiple edge is of the order $O(T^2e^{-\frac{1}{6}(\lambda-\mu)T})$. ☐

Proof of Corollary 1.4

Recall that $\tilde{\nu}_t$ denotes the distribution of the number of neighbours, ν_t the degree distribution at time t and ν the asymptotic degree distribution. Lemma 6.1 implies that $d_{TV}(\tilde{\nu}_t,\nu_t)=O(t^2e^{-\frac{1}{6}(\lambda-\mu)t})$ as $t\to\infty$. From Theorem 1.2, we know that $d_{TV}(\nu_t,\nu)=O\big(t^2e^{-\frac{1}{6}(\lambda-\mu)t}\big)$ as $t\to\infty$. Thus the triangle inequality yields the desired result. ☐

Acknowledgement

We would like to thank Thomas Rippl for pointing out to us the connection between birth and death and contour processes.

References

[AE06] Herbert Amann and Joachim Escher. *Analysis I.* Birkhäuser, Basel, 2005.

[AN72] Krishna B Athreya and Peter E Ney. *Branching processes.* Springer, Berlin, Heidelberg, New York, 1972.

[BA99] Albert-László Barabási and Réka Albert. Emergence of Scaling in Random Networks. *Science*, 286(5439):509–512, 1999.

[Bai64] Norman TJ Bailey. *The elements of stochastic processes with applications to the natural sciences.* Wiley, New York, 1964.

[BL09] Tom Britton and Mathias Lindholm. The early stage behaviour of a stochastic SIR epidemic with term-time forcing. *Journal of Applied Probability*, 46(4):975–992, 2009.

[BL10] Tom Britton and Mathias Lindholm. Dynamic random networks in dynamic populations. *Journal of Statistical Physics*, 139(3):518–535, 2010.

[BLT11] Tom Britton, Mathias Lindholm, and Tatyana Turova. A dynamic network in a dynamic population: asymptotic properties. *Journal of Applied Probability*, 48(4):1163–1178, 2011.

[BPS12] Mamadou Ba, Etienne Pardoux, and Ahmadou B Sow. Binary trees, exploration processes, and an extended Ray-Knight theorem. *Journal of Applied Probability*, 49(1):210–225, 2012.

[CJ10] Pierre Comon and Christian Jutten. *Handbook of Blind Source Separation: Independent component analysis and applications.* Academic press, 2010.

[CM77] David R Cox and Hilton D Miller. *The theory of stochastic processes*, volume 134. CRC Press, Boca Raton, Florida, 1977.

[CS12] J Theodore Cox and Rinaldo B Schinazi. A branching process for virus survival. *Journal of Applied Probability*, 49(3):888–894, 2012.

[Eis08] Bennett Eisenberg. On the expectation of the maximum of iid geometric random variables. *Statistics & Probability Letters*, 78(2):135–143, 2008.

[Fel67] William Feller. *An introduction to probability theory and its applications.* Wiley, New York, 1967.

[Ger08] Tanja Gernhard. The conditioned reconstructed process. *Journal of theoretical biology*, 253(4):769–778, 2008.

[HJV05] Patsy Haccou, Peter Jagers, and Vladimir A Vatutin. *Branching processes: variation, growth, and extinction of populations.* Cambridge University Press, Cambridge, 2005.

[Har50] Theodore Edward Harris. Some mathematical models for branching processes. Technical report, No. P-152, RAND Corporation, Santa Monica, 1950.

[Har02] Theodore Edward Harris. *The Theory of Branching Processes*. Dover Publications, Mineola, New York, 2002.

[Ken49] David G Kendall. Stochastic processes and population growth. *Journal of the Royal Statistical Society. Series B (Methodological)*, 11(2):230–282, 1949.

[Lam08] Amaury Lambert. Population dynamics and random genealogies. *Stochastic Models*, 24(S1):45–163, 2008.

[NR71] Marcel F Neuts and Sidney I Resnick. On the times of births in a linear birthprocess. *Journal of the Australian Mathematical Society*, 12(04):473–475, 1971.

[PRR13] Erol A. Peköz, Adrian Röllin, and Nathan Ross. Total variation error bounds for geometric approximation. *Bernoulli*, 19(2):610–632, 2013.

[vdH15] Remco van der Hofstad. *Random Graphs and Complex Networks. Vol. I.* Lecture Notes, 2015. `http://www.win.tue.nl/~rhofstad/NotesRGCN.pdf`

[Yan91] Nikos Yannaros. Poisson approximation for random sums of Bernoulli random variables. *Statistics & Probability Letters*, 11(2):161–165, 1991.

[Yeh95] James Yeh. *Martingales and stochastic analysis*, volume 1. World Scientific, Singapore, New Jersey, London, HongKong, 1995.